

Deep ocean sound speed characteristics passively derived from the ambient acoustic noise field

Evers, Laeslo G; Wapenaar, Kees; Heaney, Kevin D; Snellen, Mirjam

Publication date

2016

Document Version

Final published version

Citation (APA)

Evers, L. G., Wapenaar, K., Heaney, K. D., & Snellen, M. (2016). *Deep ocean sound speed characteristics passively derived from the ambient acoustic noise field*. Abstract from AGU fall meeting 2016, San Francisco, United States.

Important note

To cite this publication, please use the final published version (if applicable). Please check the document version above.

Copyright

Other than for strictly personal use, it is not permitted to download, forward or distribute the text or part of it, without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license such as Creative Commons.

Takedown policy

Please contact us and provide details if you believe this document breaches copyrights. We will remove access to the work immediately and investigate your claim.



S52A-07: Deep ocean sound speed characteristics passively derived from the ambient acoustic noise field

Friday, 16 December 2016

12:05 - 12:20

📍 Moscone South - 305

The propagation of acoustic waves in the ocean strongly depends on the temperature. Low frequency acoustic waves can penetrate the ocean down to depths where few in-situ measurements are available. It is therefore attractive to obtain a measure of the deep ocean temperature from acoustic waves. The latter is especially true if the ambient acoustic noise field can be used instead of deterministic transient signals. In this study the acoustic velocity, and hence the temperature, is derived in an interferometric approach from hydrophone array recordings. The arrays were separated by over 125 km, near Ascension Island in the Atlantic Ocean, at a depth of 800m. Furthermore, the dispersive characteristics of the deep ocean sound channel are resolved based on the retrieved lag times for different modes. In addition, it is shown how the resolution of the interferometric approach can be increased by cross correlating array beams rather than recordings from single-sensor pairs. The observed acoustic lag times between the arrays corresponds well to modeled values, based on full-wave modeling through best-known oceanic models.

Authors

[Laeslo G Evers](#) *

Royal Netherlands Meteorological Institute

[Cornelis P A Wapenaar](#)

Delft University of Technology

[Kevin D Heaney](#)

OASIS Inc.

[Mirjam Snellen](#)

Delft University of Technology

[Find Similar](#)

View Related Events

Day: [Friday, 16 December 2016](#)