

On the interplay between downwelling, deep convection and mesoscale eddies in the Labrador Sea

Georgiou, Sotiria; van der Boog, Carine; Brüggemann, Nils

Publication date

2019

Document Version

Final published version

Citation (APA)

Georgiou, S., van der Boog, C., & Brüggemann, N. (2019). On the interplay between downwelling, deep convection and mesoscale eddies in the Labrador Sea. Poster session presented at EGU General Assembly 2019 , Vienna, Austria.

Important note

To cite this publication, please use the final published version (if applicable).
Please check the document version above.

Copyright

Other than for strictly personal use, it is not permitted to download, forward or distribute the text or part of it, without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license such as Creative Commons.

Takedown policy

Please contact us and provide details if you believe this document breaches copyrights.
We will remove access to the work immediately and investigate your claim.



On the interplay between downwelling, deep convection and mesoscale eddies in the Labrador Sea

Sotiria Georgiou (1), Carine G. van der Boog (1), Nils Brüggemann (1,2), Stefanie L. Ypma (1), Julie D. Pietrzak (1), and Caroline A. Katsman (1)

(1) Delft University of Technology, Delft, Netherlands (s.georgiou@tudelft.nl), (2) Faculty of Mathematics, Informatics and Natural Sciences, University of Hamburg, Hamburg, Germany

In this study, an idealized eddy-resolving model is employed to examine the interplay between the downwelling, ocean convection and mesoscale eddies in the Labrador Sea and the spreading of dense water masses. The model output displays a basin mean net downwelling of 3.0 Sv. Our analysis confirms that the downwelling occurs near the west Greenland coast and that the eddies spawned from the boundary current play a major role in controlling the dynamics of the downwelling. The magnitude of the downwelling is positively correlated to the magnitude of the applied surface heat loss. However, we argue that this connection is indirect: the heat fluxes affect the convection properties as well as the eddy field, while the latter governs the Eulerian downwelling. A passive tracer analysis shows that dense water is transported from the interior towards the boundary, predominantly towards the Labrador coast in shallow layers and towards the Greenland coast in deeper layers. The latter transport is steered by the presence of the eddy field. The outcome that the characteristics of the downwelling in a marginal sea like the Labrador Sea depend crucially on the properties of the eddy field emphasizes that it is essential to resolve the eddies to properly represent the downwelling and overturning in the North Atlantic Ocean, and its response to changing environmental conditions.