

A Dichotomy Concerning Uniform Boundedness of Riesz Transforms on Riemannian Manifolds

Amenta, Alex; Tolomeo, Leonardo

DOI

[10.1090/proc/14730](https://doi.org/10.1090/proc/14730)

Publication date

2019

Document Version

Accepted author manuscript

Published in

American Mathematical Society. Proceedings

Citation (APA)

Amenta, A., & Tolomeo, L. (2019). A Dichotomy Concerning Uniform Boundedness of Riesz Transforms on Riemannian Manifolds. *American Mathematical Society. Proceedings*, 147(11), 4797-4803.
<https://doi.org/10.1090/proc/14730>

Important note

To cite this publication, please use the final published version (if applicable).
Please check the document version above.

Copyright

Other than for strictly personal use, it is not permitted to download, forward or distribute the text or part of it, without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license such as Creative Commons.

Takedown policy

Please contact us and provide details if you believe this document breaches copyrights.
We will remove access to the work immediately and investigate your claim.

A DICHOTOMY CONCERNING UNIFORM BOUNDEDNESS OF RIESZ TRANSFORMS ON RIEMANNIAN MANIFOLDS

ALEX AMENTA AND LEONARDO TOLOMEO

ABSTRACT. Given a sequence of complete Riemannian manifolds (M_n) of the same dimension, we construct a complete Riemannian manifold M such that for all $p \in (1, \infty)$ the L^p -norm of the Riesz transform on M dominates the L^p -norm of the Riesz transform on M_n for all n . Thus we establish the following dichotomy: given p and d , either there is a uniform L^p bound on the Riesz transform over all complete d -dimensional Riemannian manifolds, or there exists a complete Riemannian manifold with Riesz transform unbounded on L^p .

1. INTRODUCTION

Given a Riemannian manifold M , one can consider the Riesz transform $R := \nabla(-\Delta)^{\frac{1}{2}}$, where ∇ is the Riemannian gradient and Δ is the (negative) Laplace–Beltrami operator. In the Euclidean case $M = \mathbb{R}^n$, this can be identified with the vector of classical Riesz transforms (R_1, \dots, R_n) , as can be seen by writing R as a Fourier multiplier (see [12, §5.1.4]).

It is easy to show that R is bounded from $L^2(M)$ to $L^2(M; TM)$, and substantially harder to determine whether R extends to a bounded map from $L^p(M)$ to $L^p(M; TM)$ for $p \neq 2$. We let

$$R_p(M) := \sup_{\|f\|_{L^p} \leq 1} \|R(f)\|_{L^p}$$

denote the (possibly infinite) L^p -norm of the Riesz transform on M . Various conditions, often involving the heat kernel on M and its gradient, are known to imply finiteness of $R_p(M)$; see for example [2, 3, 4, 5, 6, 7, 8, 9, 13, 14]. These results usually entail finiteness of $R_p(M)$ for all $p \in (1, 2)$, or for some range of $p > 2$. On the other hand, there exist manifolds M for which $R_p(M)$ is known to be infinite for some (or all) $p > 2$: see [1, 5, 6, 7, 8, 13].

Remark 1.1. When M has finite volume we abuse notation and write $L^p(M)$ to denote the space of p -integrable functions *with mean zero*. This modification ensures that $(-\Delta)^{-1/2}$ is densely defined. When M has infinite volume, $L^p(M)$ denotes the usual Lebesgue space.

1991 *Mathematics Subject Classification*. Primary: 42B20; Secondary: 58J35, 58J65.

Key words and phrases. Riesz transform, Riemannian manifolds, Brownian motion.

The first author was supported by the VIDI subsidy 639.032.427 of the Netherlands Organisation for Scientific Research (NWO). The second author was supported by the European Research Council (grant no. 637995 “ProbDynDispEq”) and by The Maxwell Institute Graduate School in Analysis and its Applications, a Centre for Doctoral Training funded by the UK Engineering and Physical Sciences Research Council (grant EP/L016508/01), the Scottish Funding Council, Heriot-Watt University and the University of Edinburgh.

The Euclidean case is now classical: for all $p \in (1, \infty)$ there is a constant $C_p < \infty$ such that $R_p(\mathbb{R}^n) \leq C_p < \infty$ for all $n \in \mathbb{N}$ ([16]). This behaviour is expected to persist for all complete Riemannian manifolds, at least for $p < 2$. More precisely, in [9] it is conjectured that for all $p \in (1, 2)$ there exists a constant $C_p < \infty$ such that $R_p(M) \leq C_p$ for all complete Riemannian manifolds M . Such uniform bounds have been proven for all $p \in (1, \infty)$ under curvature assumptions; rather than provide an overview of the vast literature on this topic we simply point to the recent paper [10] and references therein.

One could weaken the conjecture slightly and guess that $R_p(M)$ is finite for all M , given $p \in (1, 2)$. In this article we show that this can only hold if the bound is uniform among all manifolds of a fixed dimension. This observation follows from the following dichotomy.

Theorem 1.2. *Fix $d \in \mathbb{N}$ and $p \in (1, \infty)$. Then the following dichotomy holds: either*

- *there exists a constant $C_{p,d} < \infty$ such that $R_p(M) \leq C_{p,d}$ for all complete d -dimensional Riemannian manifolds M , or*
- *there exists a complete $(d+1)$ -dimensional Riemannian manifold M such that $R_p(M) = \infty$.*

This follows from the following proposition, which we prove by an explicit construction.

Proposition 1.3. *Fix $d \geq 1$, and let $(M_n)_{n \in \mathbb{N}}$ be a sequence of complete d -dimensional Riemannian manifolds. Then there exists a complete Riemannian manifold M of dimension $d+1$ such that for all $p \in (1, \infty)$,*

$$R_p(M) \geq \sup_{n \in \mathbb{N}} R_p(M_n).$$

The main implication of Theorem 1.2 is as follows: to construct a manifold M for which $R_p(M) = \infty$ for some $p \in (1, 2)$, it suffices to construct a sequence $(M_n)_{n \in \mathbb{N}}$ of manifolds of equal dimension such that $R_p(M_n) \rightarrow \infty$ as $n \rightarrow \infty$. Thus one is led to consider lower bounds for L^p -norms of Riesz transforms. These seem not to have been considered in the literature, excluding of course the well-known computation of the L^p -norm of the Hilbert transform (the Riesz transform on \mathbb{R}) [15]. We hope that our contribution will provoke further interest in such lower bounds.

2. PRELIMINARY LEMMAS

We begin with some basic lemmas. The first says that the range of the Laplace-Beltrami operator is dense in L^p , and the second relates the Riesz transform on a manifold M with that on the M -cylinder $M \times \mathbb{R}$. These cylinders play a key role in the proof of our main theorem.

Lemma 2.1. *Let M be a complete Riemannian manifold. Then the set $S := \Delta(C_c^\infty(M))$ is dense in $L^p(M)$ for all $p \in (1, \infty)$ (recalling that we write $L^p(M)$ for the space of p -integrable mean zero functions when M has finite volume).*

Proof. Let $H \in L^p(M)$ be such that $\langle H, F \rangle = 0$ for every $F \in S$. Then $\langle H, \Delta G \rangle = 0$ for every test function G , so H is harmonic. By [17, Theorem 3], it follows that H is constant, and the result follows. \square

Lemma 2.2. *Let M be a complete Riemannian manifold. Then*

$$R_p(M \times \mathbb{R}) \geq R_p(M).$$

Proof. Consider the following modification of the Riesz transform on $M \times \mathbb{R}$:

$$\tilde{R} := \nabla_M(-\Delta_{M \times \mathbb{R}})^{-\frac{1}{2}} = \nabla_M(-\Delta_M - \partial_t^2)^{-\frac{1}{2}}.$$

This is just the projection of R onto the first summand of the tangent bundle $T(M \times \mathbb{R}) = TM \oplus T\mathbb{R}$, so we have that

$$(1) \quad \|\tilde{R}F\|_{L^p} \leq \|RF\|_{L^p}.$$

Let $F \in C_c^\infty(M \times \mathbb{R})$, and for all $\lambda > 0$ consider the function

$$F_\lambda(x, t) := \lambda^{\frac{1}{p}} F(x, \lambda t),$$

which satisfies $\|F_\lambda\|_{L^p(M \times \mathbb{R})} = \|F\|_{L^p(M \times \mathbb{R})}$. Rescaling the operator \tilde{R} in the variable t , we define

$$\tilde{R}_\lambda := \nabla_M(-\Delta_M - \lambda^2 \partial_t^2)^{-\frac{1}{2}},$$

so that

$$(2) \quad \|\tilde{R}F_\lambda\|_{L^p} = \|\tilde{R}_\lambda F\|_{L^p}.$$

Now take $f \in C_c^\infty(M) \cap D((-\Delta_M)^{-\frac{1}{2}})$ and $\rho \in C_c^\infty(\mathbb{R})$ such that $\|\rho\|_{L^p(\mathbb{R})} = 1$, and consider the function $F(x, t) = f(x)\rho(t)$. Since Δ_M and ∂_t^2 commute, and the function

$$G_\lambda(x, y) = \left(\frac{x}{x + \lambda^2 y} \right)^{\frac{1}{2}}$$

is bounded by 1 for $(x, y) > 0$, and $G_\lambda \rightarrow 1$ pointwise as $\lambda \rightarrow 0$, we have

$$\lim_{\lambda \rightarrow 0} (-\Delta_M - \lambda^2 \partial_t^2)^{-\frac{1}{2}} F = \lim_{\lambda \rightarrow 0} G_\lambda(-\Delta_M, -\partial_t^2)(-\Delta_M)^{-\frac{1}{2}} f \otimes \rho = (-\Delta_M)^{-\frac{1}{2}} f \otimes \rho$$

in L^2 , and thus also as distributions. Therefore $\tilde{R}_\lambda F \rightarrow Rf \otimes \rho$ as distributions, and so

$$\liminf_{\lambda \rightarrow 0} \|\tilde{R}_\lambda F\|_{L^p(M \times \mathbb{R})} \geq \|Rf \otimes \rho\|_{L^p(M \times \mathbb{R})} = \|Rf\|_{L^p(M)}.$$

Combining this with (2) and (1), and the fact that $C_c^\infty(M) \cap D((-\Delta_M)^{-\frac{1}{2}})$ is dense in $L^p(M)$,¹ yields $R_p(M \times \mathbb{R}) \geq R_p(M)$. \square


3. PROOF OF THE MAIN THEOREM

In this section we carry out the construction that proves Proposition 1.3, which implies Theorem 1.2.

Consider a sequence $(M_n)_{n \in \mathbb{N}}$ of complete d -dimensional Riemannian manifolds. We will connect the M_n -cylinders $(M_n \times \mathbb{R})_{n \in \mathbb{N}}$ along a \mathbb{T}^d -cylinder $\mathbb{T}^d \times \mathbb{R}$ as follows.² For each $n \in \mathbb{N}$ fix a coordinate chart $U_n \subset M_n \times (-1/2, 1/2)$ and a small ball $B_n \subset U_n$. Similarly, for each $n \in \mathbb{N}$ choose a small coordinate chart $U'_n \subset \mathbb{T}^n \times \mathbb{R}$ such that the charts $(U'_n)_{n \in \mathbb{N}}$ are pairwise disjoint, and a small ball $B'_n \subset U'_n$. For each $n \in \mathbb{N}$, glue the manifold $(M_n \times \mathbb{R}) \setminus B_n$ to $(\mathbb{T}^n \times \mathbb{R}) \setminus B'_n$ along the boundaries

¹This follows from the inclusion $D((-\Delta_M)^{-\frac{1}{2}}) \supseteq D((-\Delta_M)^{-1}) \supseteq \Delta_M(C_c^\infty(M))$, which is dense by Lemma 2.1. See also [11, Lemma 2.2]. Again, recall that $L^p(M)$ denotes the corresponding space of mean zero functions when M has finite volume.

²Of course, one could connect the M_n -cylinders to each other directly, without needing the \mathbb{T}^d -cylinder. This would work just as well.

FIGURE 1. Construction of M from $(M_n)_{n \in \mathbb{N}}$.

of B_n and B'_n ; this is possible since both these balls are ‘Euclidean’ balls sitting inside coordinate charts. This results in a C^0 -Riemannian manifold (M, g') , which is C^∞ away from the set $\Sigma = \cup_n \partial B_n$ on which we glued the manifolds together. Mollify the metric to get a C^∞ -Riemannian manifold (M, g) such that $g = g'$ away from the ε -neighbourhood of Σ for some very small ε . An artist’s impression of this construction, with $M_n = S^1$ for each n , is shown in Figure 1.

For each $n \in \mathbb{N}$ we have an inclusion map

$$i_n: M_n \times (1, \infty) \rightarrow M$$

which is an isometry. From here on we fix n and just write $i = i_n$. Functions on M can be pulled back to $M_n \times (1, \infty)$; the pullback map is denoted i^* , so that for $f: M \rightarrow \mathbb{R}$ the function $i^*f: M_n \times (1, \infty) \rightarrow \mathbb{R}$ is defined by

$$i^*f(x, t) = f(i(x, t)).$$

On the other hand, for $g: M_n \times (1, \infty) \rightarrow \mathbb{R}$ we can define a pushforward $i_*g: M \rightarrow \mathbb{R}$ by setting $i_*g(i(x, t)) := g(x, t)$ on $i(M_n \times (1, \infty))$ and extending by zero to the rest of M . For a function $g: M_n \times \mathbb{R} \rightarrow \mathbb{R}$ and for $s \in \mathbb{R}$ we let $\tau_s g: M_n \times \mathbb{R} \rightarrow \mathbb{R}$ be the translated function $\tau_s g(x, t) := g(x, t - s)$. Similarly if $g: M_n \times (1, \infty) \rightarrow \mathbb{R}$ we can define $\tau_s g: M_n \times (1 + s, \infty) \rightarrow \mathbb{R}$. These concepts apply equally well to vector fields in place of functions.

We will need the following lemma, which relates the heat flow on $M_n \times \mathbb{R}$ to the one on M .

Lemma 3.1. *Let $F: M_n \times \mathbb{R} \rightarrow \mathbb{R}$ be smooth and compactly supported, and fix $\sigma > 0$. Then for every $(x, t) \in M_n \times \mathbb{R}$,*

$$\lim_{s \rightarrow +\infty} (e^{\sigma \Delta_M} i_* \tau_s F)(i(x, t + s)) = (e^{\sigma \Delta_{M_n \times \mathbb{R}}} F)(x, t).$$

Proof. Let $W_{x,t}(\sigma)$ be a Brownian motion on $M_n \times \mathbb{R}$ at time σ starting from the point (x, t) . Since the generator $\frac{1}{2}\Delta_{M_n \times \mathbb{R}}$ satisfies $\frac{1}{2}i_* \circ \Delta_{M \times \mathbb{R}}|_{i(M_n \times (1, +\infty))} = \frac{1}{2}\Delta_M|_{i(M_n \times (1, +\infty))}$, defining the stopping time

$$T(x, t) := \inf \{s : W_{x,t}(s) \in M_n \times (-\infty, 1)\},$$

we have that $i(W_{x,t}(\sigma))$ is a Brownian motion on M for $\sigma < T(x, t)$. Therefore there exists a Brownian motion $\tilde{W}_{i(x,t)}(\sigma)$ on M such that $\tilde{W}(\sigma) = i(W(\sigma))$ for $\sigma < T$; if \bar{W} is a Brownian motion on M , we can take for example

$$\tilde{W}_{i(x,t)}(\sigma) = \begin{cases} i(W_{x,t}(\sigma)) & \text{if } \sigma < T, \\ \bar{W}_{i(W_{x,t}(T))}(\sigma - T) & \text{if } \sigma \geq T. \end{cases}$$

We have that

$$\begin{aligned} & (e^{\sigma\Delta_M} i_* \tau_s F)(i(x, t+s)) \\ &= \mathbb{E}[(i_* \tau_s F)(\tilde{W}_{i(x,t+s)}(2\sigma))] \\ &= \mathbb{E}[(i_* \tau_s F)(\tilde{W}_{i(x,t+s)}(2\sigma))\mathbb{1}_{2\sigma < T}] + \mathbb{E}[(i_* \tau_s F)(\tilde{W}_{i(x,t+s)}(2\sigma))\mathbb{1}_{2\sigma \geq T}] \\ &= \mathbb{E}[(\tau_s F)(W_{x,t+s}(2\sigma))\mathbb{1}_{2\sigma < T}] + \mathbb{E}[(i_* \tau_s F)(\tilde{W}_{i(x,t+s)}(2\sigma))\mathbb{1}_{2\sigma \geq T}] \\ &= \mathbb{E}[(\tau_s F)(W_{x,t+s}(2\sigma))] \\ &\quad - \mathbb{E}[(\tau_s F)(W_{x,t+s}(2\sigma))\mathbb{1}_{2\sigma \geq T}] + \mathbb{E}[(i_* \tau_s F)(\tilde{W}_{i(x,t+s)}(2\sigma))\mathbb{1}_{2\sigma \geq T}] \\ &= (e^{\sigma\Delta_{M_n \times \mathbb{R}}} \tau_s F)(x, t+s) \\ &\quad - \mathbb{E}[(\tau_s F)(W_{x,t+s}(2\sigma))\mathbb{1}_{2\sigma \geq T}] + \mathbb{E}[(i_* \tau_s F)(\tilde{W}_{i(x,t+s)}(2\sigma))\mathbb{1}_{2\sigma \geq T}]. \end{aligned}$$

Therefore

$$|(e^{\sigma\Delta_M} i_* \tau_s F)(i(x, t+s)) - (e^{\sigma\Delta_{M_n \times \mathbb{R}}} \tau_s F)(x, t+s)| \leq 2\|F\|_{L^\infty} \mathbb{P}(T(x, t+s) \leq 2\sigma).$$

Since $\Delta_{M_n \times \mathbb{R}}$ is translation invariant in the \mathbb{R} coordinate, we have that

$$\begin{aligned} \mathbb{P}(T(x, t+s) \leq 2\sigma) &\leq \mathbb{P}(\{W_{x,t+s}(\sigma') \in M_n \times (-\infty, 1) \text{ for some } \sigma' \leq 2\sigma + 1\}) \\ &= \mathbb{P}(\{W_{x,t}(\sigma') \in M_n \times (-\infty, 1-s) \text{ for some } \sigma' \leq 2\sigma + 1\}) \end{aligned}$$

and by continuity of $W_{x,t}(\cdot)$, this tends to 0 as $s \rightarrow \infty$. Thus we find that

$$\lim_{s \rightarrow +\infty} \left((e^{\sigma\Delta_M} i_* \tau_s F)(i(x, t+s)) - (e^{\sigma\Delta_{M_n \times \mathbb{R}}} \tau_s F)(x, t+s) \right) = 0.$$

The conclusion follows from translation invariance of $\Delta_{M_n \times \mathbb{R}}$ in \mathbb{R} . \square

We return to the proof of Proposition 1.3. Fix $\varepsilon > 0$, and choose $F = \Delta_{M_n \times \mathbb{R}} H$ for some $H \in C_c^\infty(M_n \times \mathbb{R})$ with $\|F\|_{L^p} = 1$ such that

$$\|R_{M_n \times \mathbb{R}} F\|_{L^p} \geq (R_p(M_n) - \varepsilon) \wedge \varepsilon^{-1}.$$

Such a function exists by Lemmas 2.1 and 2.2. We claim that

$$(3) \quad \lim_{s \rightarrow +\infty} \tau_{-s} i^* R_M(i_* \tau_s F) = R_{M_n \times \mathbb{R}} F$$

as distributions. Assuming (3) for the moment, we have

$$\begin{aligned} \limsup_{s \rightarrow \infty} \|R_M(i_* \tau_s F)\|_{L^p(M)} &\geq \limsup_{s \rightarrow \infty} \|i^* R_M(i_* \tau_s F)\|_{L^p(M_n \times \mathbb{R})} \\ &= \limsup_{s \rightarrow \infty} \|\tau_{-s} i^* R_M(i_* \tau_s F)\|_{L^p(M_n \times \mathbb{R})} \\ &\geq \|R_{M_n \times \mathbb{R}} F\|_{L^p(M_n \times \mathbb{R})} \geq R_p(M_n) - \varepsilon, \end{aligned}$$

while for all $s \in \mathbb{R}$

$$\|i_*\tau_s F\|_{L^p(M)} \leq \|\tau_s F\|_{L^p(M_n \times \mathbb{R})} = \|F\|_{L^p(M_n \times \mathbb{R})} \leq 1.$$

The result follows, so it remains to prove (3).

For s sufficiently large, we have that

$$i_*\tau_s F = i_*\tau_s(\Delta_{M_n \times \mathbb{R}} H) = i_*(\Delta_{M_n \times \mathbb{R}} \tau_s H) = \Delta_M i_*\tau_s H,$$

therefore $i_*\tau_s F \in D(\Delta_M^{-1}) \subseteq D((-\Delta_M)^{-\frac{1}{2}})$, and hence

$$R(i_*\tau_s F) = \nabla \left((-\Delta_M)^{-\frac{1}{2}} i_*\tau_s F \right)$$

as a distribution. To test the distributional convergence, let X be a smooth compactly supported vector field in $M_n \times \mathbb{R}$. For large s we have that

$$\begin{aligned} \langle \tau_{-s} i^* R_M(i_*\tau_s F), X \rangle &= \langle R_M(i_*\tau_s F), i_*\tau_s X \rangle \\ &= \left\langle (-\Delta_M)^{-\frac{1}{2}} i_*\tau_s F, \operatorname{div}(i_*\tau_s X) \right\rangle \\ &= \left\langle (-\Delta_M)^{-\frac{1}{2}} i_*\tau_s F, i_*\tau_s \operatorname{div}(X) \right\rangle. \end{aligned}$$

Therefore it is enough to show that for every $G \in C_c^\infty(M_n \times \mathbb{R})$,

$$(4) \quad \lim_{s \rightarrow \infty} \left\langle (-\Delta_M)^{-\frac{1}{2}} i_*\tau_s F, i_*\tau_s G \right\rangle = \left\langle (-\Delta_{M_n \times \mathbb{R}})^{-\frac{1}{2}} F, G \right\rangle.$$

By the well-known formula

$$(-\Delta)^{-\frac{1}{2}} = \pi^{-\frac{1}{2}} \int_0^{+\infty} \sigma^{-\frac{1}{2}} e^{\sigma \Delta} d\sigma,$$

(4) is equivalent to showing that

$$(5) \quad \lim_{s \rightarrow \infty} \int_0^{+\infty} \sigma^{-\frac{1}{2}} \langle e^{\sigma \Delta_M} i_*\tau_s F, i_*\tau_s G \rangle d\sigma = \int_0^{+\infty} \sigma^{-\frac{1}{2}} \langle e^{\sigma \Delta_{M_n \times \mathbb{R}}} F, G \rangle d\sigma.$$

Note that

$$\left| \sigma^{-\frac{1}{2}} \langle e^{\sigma \Delta_M} i_*\tau_s F, i_*\tau_s G \rangle \right| \leq \sigma^{-\frac{1}{2}} \|i_*\tau_s F\|_{L^2} \|i_*\tau_s G\|_{L^2} \leq \sigma^{-\frac{1}{2}} \|F\|_{L^2} \|G\|_{L^2}$$

and

$$\left| \sigma^{-\frac{1}{2}} \langle e^{\sigma \Delta_M} i_*\tau_s F, i_*\tau_s G \rangle \right| = \left| \sigma^{-\frac{3}{2}} \langle e^{\sigma \Delta_M} \sigma \Delta_M i_*\tau_s H, i_*\tau_s G \rangle \right| \lesssim \sigma^{-\frac{3}{2}} \|H\|_{L^2} \|G\|_{L^2}.$$

Since the function $\min(\sigma^{-\frac{1}{2}}, \sigma^{-\frac{3}{2}})$ is integrable, by dominated convergence (5) will be proved if we show

$$(6) \quad \lim_{s \rightarrow \infty} \langle e^{\sigma \Delta_M} i_*\tau_s F, i_*\tau_s G \rangle = \langle e^{\sigma \Delta_{M_n \times \mathbb{R}}} F, G \rangle$$

for every $\sigma > 0$. We show (6) by writing

$$\begin{aligned} \lim_{s \rightarrow \infty} \langle e^{\sigma \Delta_M} i_*\tau_s F, i_*\tau_s G \rangle &= \lim_{s \rightarrow \infty} \langle \tau_{-s} i^* e^{\sigma \Delta_M} i_*\tau_s F, G \rangle \\ &= \lim_{s \rightarrow \infty} \int_{1-s}^{+\infty} \int_{M_n} (e^{\sigma \Delta_M} i_*\tau_s F)(i(x, t+s)) G(x, t) dx dt \\ &= \int_{\mathbb{R}} \int_{M_n} (e^{\sigma \Delta_{M_n \times \mathbb{R}}} F)(x, t) G(x, t) dx dt \\ &= \langle e^{\sigma \Delta_{M_n \times \mathbb{R}}} F, G \rangle, \end{aligned}$$

using Lemma 3.1 and dominated convergence (by $\|F\|_{L^\infty} |G(x, t)|$). This completes the proof of Proposition 1.3, and hence establishes Theorem 1.2.

REFERENCES

- [1] A. Amenta. New Riemannian manifolds with L^p -unbounded Riesz transform for $p > 2$. arXiv:1707.09781, July 2017.
- [2] P. Auscher and T. Coulhon. Riesz transform on manifolds and Poincaré inequalities. *Ann. Sc. Norm. Super. Pisa Cl. Sci. (5)*, 4(3):531–555, 2005.
- [3] P. Auscher, T. Coulhon, X. T. Duong, and S. Hofmann. Riesz transform on manifolds and heat kernel regularity. *Ann. Sci. École Norm. Sup. (4)*, 37(6):911–957, 2004.
- [4] F. Bernicot and D. Frey. Riesz transforms through reverse Hölder and Poincaré inequalities. *Math. Z.*, 284(3-4):791–826, 2016.
- [5] G. Carron. Riesz transform on manifolds with quadratic curvature decay. *Rev. Mat. Iberoam.*, 33(3):749–788, 2017.
- [6] G. Carron, T. Coulhon, and A. Hassell. Riesz transform and L^p -cohomology for manifolds with Euclidean ends. *Duke Math. J.*, 133(1):59–93, 2006.
- [7] L. Chen, T. Coulhon, J. Feneuil, and E. Russ. Riesz transform for $1 \leq p \leq 2$ without Gaussian heat kernel bound. *J. Geom. Anal.*, 27(2):1489–1514, 2017.
- [8] T. Coulhon and X. T. Duong. Riesz transforms for $1 \leq p \leq 2$. *Trans. Amer. Math. Soc.*, 351(3):1151–1169, 1999.
- [9] T. Coulhon and X. T. Duong. Riesz transform and related inequalities on noncompact Riemannian manifolds. *Comm. Pure Appl. Math.*, 56(12):1728–1751, 2003.
- [10] K. Dahmani, K. Domelevo, and S. Petermichl. Dimensionless L^p estimates for the Riesz vector on manifolds. arXiv:1802.00366, February 2018.
- [11] Baptiste Devyver. A perturbation result for the Riesz transform. *Ann. Sc. Norm. Super. Pisa Cl. Sci. (5)*, 14(3):937–964, 2015.
- [12] L. Grafakos. *Classical Fourier analysis*, volume 249 of *Graduate Texts in Mathematics*. Springer, New York, third edition, 2014.
- [13] H.-Q. Li. La transformation de Riesz sur les variétés coniques. *J. Funct. Anal.*, 168(1):145–238, 1999.
- [14] H.-Q. Li and J.-X. Zhu. A note on “Riesz transform for $1 \leq p \leq 2$ without Gaussian heat kernel bound”. *J. Geom. Anal.*, Jun 2017. Published online.
- [15] S. K. Pichorides. On the best values of the constants in the theorems of M. Riesz, Zygmund and Kolmogorov. *Studia Math.*, 44:165–179, 1972.
- [16] E. M. Stein. Some results in harmonic analysis in \mathbf{R}^n , for $n \rightarrow \infty$. *Bull. Amer. Math. Soc. (N.S.)*, 9(1):71–73, 1983.
- [17] S. T. Yau. Some function-theoretic properties of complete Riemannian manifold and their applications to geometry. *Indiana Univ. Math. J.*, 25(7):659–670, 1976.

DELFT INSTITUTE OF APPLIED MATHEMATICS, DELFT UNIVERSITY OF TECHNOLOGY, P.O. BOX 5031, 2600 GA DELFT, THE NETHERLANDS
E-mail address: `amenta@fastmail.fm`

SCHOOL OF MATHEMATICS, THE UNIVERSITY OF EDINBURGH AND MAXWELL INSTITUTE FOR MATHEMATICAL SCIENCES, JAMES CLERK MAXWELL BUILDING, RM 5210 THE KING’S BUILDINGS, PETER GUTHRIE TAIT ROAD, EDINBURGH, EH9 3FD, UNITED KINGDOM
E-mail address: `L.Tolomeo@sms.ed.ac.uk`