



Delft University of Technology

Document Version

Final published version

Citation (APA)

Tsalidis, G. A. (2022). Human Health and Ecosystem Quality Benefits with Life Cycle Assessment Due to Fungicides Elimination in Agriculture. *Sustainability (Switzerland)*, 14(2), Article 846. <https://doi.org/10.3390/su14020846>

Important note

To cite this publication, please use the final published version (if applicable).
Please check the document version above.

Copyright

In case the licence states "Dutch Copyright Act (Article 25fa)", this publication was made available Green Open Access via the TU Delft Institutional Repository pursuant to Dutch Copyright Act (Article 25fa, the Taverne amendment). This provision does not affect copyright ownership.

Unless copyright is transferred by contract or statute, it remains with the copyright holder.

Sharing and reuse

Other than for strictly personal use, it is not permitted to download, forward or distribute the text or part of it, without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license such as Creative Commons.

Takedown policy

Please contact us and provide details if you believe this document breaches copyrights.
We will remove access to the work immediately and investigate your claim.

This work is downloaded from Delft University of Technology.



Review

Human Health and Ecosystem Quality Benefits with Life Cycle Assessment Due to Fungicides Elimination in Agriculture

Georgios Archimidis Tsalidis ^{1,2}

¹ Biotechnology Department, Applied Sciences Faculty, Delft University of Technology, 2629 HZ Delft, The Netherlands; g.a.tsalidis@tudelft.nl or gatsalidis@athenarc.gr

² Athena—Research and Innovation Center in Information, Communication and Knowledge Technologies, 67100 Xanthi, Greece

Abstract: Industrial agriculture results in environmental burdens due to the overuse of fertilizers and pesticides. Fungicides is a class of pesticides whose application contributes (among others) to human toxicity and ecotoxicity. The European Union aims to increase organic agriculture. For this reason, this work aims to analyze climate change, freshwater ecotoxicity, terrestrial ecotoxicity, human toxicity, (terrestrial) acidification, and freshwater eutrophication impacts of fungicides and calculate expected benefits to human health (per European citizen) and ecosystem quality (terrestrial) with life cycle assessment (LCA) during crop production. The Scopus database was searched for LCA studies that considered the application of fungicides to specific crops. The analysis shows how many systemic and contact fungicides were considered by LCA studies and what was the applied dosage. Furthermore, it shows that fungicides highly contribute to freshwater ecotoxicity, terrestrial ecotoxicity, human toxicity, and freshwater eutrophication for fruits and vegetables, but to a low extent compared to all considered environmental impacts in the case of cereals and rapeseed. Expected benefits to human health and ecosystem quality after fungicides elimination are greater for fruits and vegetables, ranging between 0 to 47 min per European citizen in a year and 0 to 90 species per year, respectively.



Citation: Tsalidis, G.A. Human Health and Ecosystem Quality Benefits with Life Cycle Assessment Due to Fungicides Elimination in Agriculture. *Sustainability* **2022**, *14*, 846. <https://doi.org/10.3390/su14020846>

Academic Editor: Idiano D'Adamo

Received: 21 December 2021

Accepted: 11 January 2022

Published: 12 January 2022

Publisher's Note: MDPI stays neutral with regard to jurisdictional claims in published maps and institutional affiliations.



Copyright: © 2022 by the author. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).

1. Introduction

The most environmentally damaging form of human consumption is eating [1] because industrial agriculture results in environmental burdens due to the overuse of machinery, fertilizers, insecticides, herbicides, and fungicides [2]. The latter are toxic to humans and the environment [3]. This study reviews the contribution to toxicity of fungicides in fruits, vegetables, cereals, and rapeseed production to show the benefits for human health and ecosystem quality after their elimination.

The primary goals of agriculture are the production of nutritional food for humans and feed for animals, and the increased economic development of stakeholders. In addition, the UN 2030 Agenda on Sustainable Development also identified bioeconomy as being in line with goals for energy efficiency (Goal 7) [4]. These goals resulted in a significant growth of the bioeconomy, and the number of publications about bioeconomy increased from 1000 in 2017 to approximately 3500 in 2021 [5]. Furthermore, during the period 2021 to 2030, global agricultural production is projected to increase by 1.4% per year [6]. Maize, rice, wheat, oilseeds, and oil products are major commodities globally [6]. Due to the introduction of synthetic pesticides between 1960 and 1989, pest control and agricultural output have significantly improved [7].

Intensive, industrial agricultural practices employ large quantities of inputs, such as fuels, electricity, water for irrigation, fertilizers, and pesticides. Such intensive agricultural

practices are held responsible for increased energy use and accompanied elevated emissions of greenhouse gases [8], and adverse effects on human health and ecosystem quality [3]. In contrast, in organic agricultural practice, the use of fertilizers is significantly reduced while the consumption of pesticides is avoided [9].

The term pesticide covers a wide range of compounds, including insecticides, fungicides, herbicides, etc. The benefits of pesticides are the consequences of their effects: increased productivity, protection of crop losses, vector disease control, and quality of food [7]. These effects occur due to active ingredients in pesticides, which are chemicals that kill, control, or repel pests. The World Health Organization [10] and a study [11] focusing on toxicity stated that insecticides tend to be more toxic to humans than fungicides and herbicides by one and four orders of magnitude, respectively. However, three active fungicides ingredients—(1) tebuconazole, (2) epoxiconazole, and (3) prochloraz—were more toxic than herbicides and insecticides to mitochondrial activities, membrane degradations, and caspases 3/7 activities [12].

Fungicides are the class of pesticides that target fungi. Fungicides are classified as systemic and contact. Systemic fungicide can be defined as a fungicide that, when sprayed on top surfaces of the plant, is translocated to the plant tissues, transferring its toxicity to the targeted fungus [13]. Examples of systemic fungicides are Benomyl, Cryoconazol, and Imazalil. Contact fungicides remain on the plant surface after application and do not penetrate plant tissues. These fungicides are typically employed to control foliar diseases. A contact fungicide must be present on the plant surface before the disease penetrates the tissue [14]. Examples of contact fungicides are Mancozeb and Thiram. Pre-harvest fungicide use is greatly limited or absent in organic agroecosystems [15]. According to studies [16–18], approximately 55% of applied fungicide is found in air, freshwater, and soil while the rest remains in or on the crop, depending on fungicide class.

Stakeholders from corporate, academic, civil society, and governmental sectors highlighted a shift to organic agriculture as a sustainable solution to the negative effects of industrial agriculture [19]. In particular, European Union (EU) strategies regarding agricultural development encourage eco-friendly practices [20]. Europe's Farm to Fork (F2F) strategy aims to boost the development of its organic farming area to 25% of total farmland by 2030. However, a recent study [21] concluded that increasing organic cultivation with the F2F strategy may result in less sustainable food systems because EU law strictly prohibits the use of genetically modified organisms in organic cultivation, which can result in failing to achieve hunger reduction (Sustainable Development Goal 2).

Life cycle assessment (LCA) is a tool to assess environmental sustainability during the entire life cycle of a product and it is standardized by ISO 14040 [22] and 14044 [23]. Energy and material flows, and environmental releases are quantified and converted to environmental impacts during the life cycle inventory and life cycle impact assessment phases, respectively. Environmental impacts exist at midpoint and endpoint levels. Midpoint indicators focus on single environmental impacts, such as climate change. The ReCiPe is an environmental impact model which calculates midpoint indicators and converts them into endpoint indicators based on an individualist, hierarchist, or egalitarian perspective. ReCiPe's endpoint indicators are human health, ecosystem quality, and resource scarcity [24].

A separate impact assessment modeling step is required for fungicides because reference life cycle inventory datasets of fungicides reflect only production-related impacts, not impacts that occur after application [3]. This review aims to analyze the environmental impacts of fungicides due to their production and application, and calculate expected benefits to human health and ecosystem quality due to their elimination based on ReCiPe's midpoint to endpoint conversion factors.

2. Materials and Methods

This study considered 39 papers. They are the results of searching in the Scopus database with certain combinations of keywords with the "AND" Boolean operator. Figure 1 shows all the steps followed to identify the considered LCA studies. To summarize,

considered studies needed to apply LCA in agricultural products, mention fungicides or at least include them in the class of pesticides during the contribution analysis of life cycle interpretation stage, and calculate toxicity. In 2009, van Zelm et al. [25] published their work which calculated freshwater ecotoxicological effect factors for 397 pesticides. Therefore, we considered this study the starting point for this research. All considered studies can be found in the Supplementary Materials.




Figure 1. Steps followed for review and inclusion/exclusion criteria.

Analysis of reviewed midpoint impact values to design box plots was performed with MS Excel. Furthermore, outliers were analyzed with Grubbs' test with a significance level of 0.05 (two-sided) in the GraphPad Outlier calculator.

Midpoint impacts were converted to endpoint damages to human health and ecosystem quality (terrestrial) with the ReCiPe environmental impact model [26]. Midpoint to endpoint conversion factors were used to convert average midpoint results (per type of crop) to human health and ecosystem quality (terrestrial) endpoints based on the individualist, hierarchist, and egalitarian perspectives, and show the effect of fungicide elimination on a European level. The following formula was used:

$$\text{Endpoint value}_i = \sum \text{average midpoint value}_{j,k} \times \text{contribution to midpoint}_{\text{fungicide}} \times \text{contribution to endpoint}_{\text{perspective}} \times \text{yield}_k \quad (1)$$

where i stands for endpoint human health or ecosystem quality (terrestrial), j stands for midpoint indicator, and k stands for the type of crop. In the case of human health, the endpoint value was divided per European population in 2019 (513.5 million people) to normalize it per person in Europe. In the case of ecosystem quality (terrestrial), the endpoint value is presented to show species loss in one year.

3. Results and Discussion

3.1. Active Ingredient

Various types of fungicides and active ingredients are used in agriculture. The same systemic or contact fungicide is used in more than one fruit, vegetable, and cereal and rapeseed crop. Table 1 shows how many systemic and contact fungicides are applied per agricultural product. Systemic fungicides are applied more than contact fungicides in terms of different active ingredients. LCA studies that focus on fruits use a smaller number of fungicides than vegetables, cereals, and rapeseed. This does not necessarily mean that the quantity of active ingredients (dosage) is smaller in fruits than vegetables, cereals, or

rapeseed. In contrast, vegetables employ the largest variety of fungicides, with systemic pesticides dominating, especially azoxystrobin [27–29] and boscalid [27,30,31]. For both fruits and vegetables, the most common contact fungicide is mancozeb [8,11,19,28,29,32]. Lastly, LCA studies of cereals and rapeseed considered almost only systemic fungicides, with azoxystrobin [33,34], propiconazole [34,35], and epoxiconazole dominating the list [34,36]. A detailed version of Table 1 can be found in the Supplementary Materials.

Table 1. Applied fungicides per agricultural product.

Product	Systemic Fungicides (Number)	Contact Fungicides (Number)
Fruits	6	4
Vegetables	16	8
Cereals and rapeseed	10	1

3.2. Dosage

The dosage of fungicides per functional unit is presented based on their active ingredients. These active ingredients are the cause of environmental impacts due to their release to the soil, water, and/or air, and residues in agricultural product. Figure 2 shows the dosage in mass per mass of fruit, vegetable, or cereal and rapeseed. LCA studies of fruits, cereals, and rapeseed consider the highest dosage of fungicides, while vegetables' dosage is much lower. For fruits, grape cultivation in Spain presents the highest dosage of copper-based fungicides [37]. For vegetables, the authors do not disclose what are the fungicides or the active ingredients for cucumber [38] and bean [39] cultivations in Iran and Greece, respectively. Regarding cereals and rapeseed, fungicides used in cotton cultivation in Mali [40] are outliers among reviewed studies because furathiocarb, metalaxyl, and chlorothalonil are supplied almost one order of magnitude more than the second-largest applied fungicide. Rice cultivation in China has the second-largest value because applied fungicides are presented with insecticides and herbicides as pesticides [41]. Lastly, climate and soil type may have an effect on the dosage of applied fungicides, but these two parameters are rarely mentioned in LCA studies.




Figure 2. Box plot with median and outlier of active ingredients dosage (of fungicide) per 1 kg of product.

3.3. Fungicides Contribution to Environmental Impacts

Active ingredients of fungicides contribute to various environmental impacts depending on their constituents and release in terms of mass, and release in terms of environmental medium. Figures 3–5 show the extent to which active ingredients contribute to climate change, freshwater ecotoxicity, terrestrial ecotoxicity, human toxicity [3], (terrestrial) acidification [42], and freshwater eutrophication [43]. Among these environmental impacts, only

climate change is not affected during fungicide application but derives from the production stage due to greenhouse gases emissions. The other impacts occur during fungicides application in the cultivation stage.




Figure 3. Box plot with median and outliers of fungicides contribution to environmental impacts of 1 kg of fruits.




Figure 4. Box plot with median and outliers of fungicides contribution to environmental impacts of 1 kg of vegetables.

The most commonly assessed food product in reviewed LCA studies was grape, and various fungicides were applied during its cultivation. Regarding fruits, terrestrial ecotoxicity, freshwater ecotoxicity, and freshwater eutrophication depend on fungicides based on the reviewed studies (see Figure 3). The range of contribution is large for these environmental impacts and different fungicides are applied. For instance, contribution to terrestrial ecotoxicity increased from 8% to 96% for grape production in South Africa [32] and cherry production in Portugal [44] due to an order of magnitude difference in fungicide dosage during cultivation and the fact that fungicides are grouped with insecticides (in the

life cycle interpretation stage), respectively. Furthermore, Beauchet et al. [45] presented a 40% contribution to terrestrial ecotoxicity and a 60% contribution to freshwater ecotoxicity and freshwater eutrophication in grape production in France, but these authors only mentioned that they considered a synthetic fungicide and grouped it with insecticides. In total, eight studies assessed freshwater ecotoxicity, which varied between 1% and 90% contribution. The contribution of 1% regards grape production in Iran and freshwater ecotoxicity is dominated by the use of poultry manure [46], while the name of the applied fungicide is not mentioned. In contrast, large quantities of applied fungicides (i.e., Fosetyl-Al and Imazalil) in orange production in Spain resulted in a 90% contribution to freshwater ecotoxicity [11]. Applied fungicides in fruit production contribute to a small extent to climate change [32,44–48], terrestrial acidification [32,44,45,47,48], and human toxicity [11,46–48] due to low characterization factors of fungicides for these impacts and other contributing environmental releases.




Figure 5. Box plot with median and outliers of fungicides contribution to environmental impacts of 1 kg of cereals and seeds.

Finally, the only distinction between systemic and contact fungicides can be observed for the contact fungicide mancozeb, which is a key contributor to freshwater ecotoxicity and freshwater eutrophication based on [11,32].

The most commonly assessed vegetable product in reviewed LCA studies is tomato. Concerning vegetables, human toxicity, freshwater ecotoxicity, terrestrial ecotoxicity, and freshwater eutrophication depend on fungicides but show a large range (see Figure 4). High contribution to toxicity is found in one study [49] for tomato production in China due to the application of iprodione (>58% regarding human toxicity) and chlorothalonil (90% regarding aquatic ecotoxicity). Similarly, tomato [50,51] and endive [31] production also show a high contribution to freshwater ecotoxicity, from 40% to 66%, respectively, but only Yelboga [51] distinguished field emissions among various pesticides and reported organo-phosphorus and triazine compounds (i.e., a fungicide) as having the highest and lowest effects, respectively. The range of contribution of freshwater eutrophication varied from 2% to 58%, but only three studies [31,36,50] were identified and all authors considered different fungicides and vegetables. Human toxicity varied between 8% and 89%. The 8% contribution is derived from potato production in Spain, where a mix of fungicides rich in mancozeb and chlorothalonil is applied [36]. On the other hand, the 89% contribution is derived from tomato production in China, where iprodione and chlorothalonil were applied [49]. The range of terrestrial ecotoxicity is smaller than freshwater ecotoxicity,

freshwater eutrophication, and human toxicity, and varied between 4% and 40%. Two studies presented a 40% contribution, but only Yelboğa [51] stated that applied fungicides acetamide anillide, phthalimide, and organo-phosphorus compounds contribute 4.4%, 4.8%, and 14.8%, respectively. Applied fungicides in vegetables production contribute to a small extent to climate change [19,29,30,36,38,49] and terrestrial acidification [28,30,31,36,38].

In general, most applied fungicides are systemic, with boscalid and azoxystrobin being the most common, while the most common contact fungicide is mancozeb. Boscalid contributes to freshwater ecotoxicity and human toxicity, whereas mancozeb contributes to climate change due to its production stage, and terrestrial acidification. Finally, it should be noted that even though the EU expects an improved environmental performance due to increasing organic production, Pedretti et al. [30] reported no statistical difference due to pesticides application between organic and conventional production of spinach leaf.

The most commonly assessed cereal product in reviewed LCA studies was rice. Wheat and rapeseed were also popular choices among researchers. Figure 5 shows the contribution of fungicides to environmental impacts due to cereals and rapeseed production. The contribution of fungicides to climate change, human toxicity, freshwater ecotoxicity, terrestrial ecotoxicity, and freshwater eutrophication was very low compared to fruits and vegetables. Contribution ranged from 0% to 6% for most impacts, except for freshwater ecotoxicity. In freshwater ecotoxicity, the application of systemic fungicides (such as epoxiconazole, propiconazole, and azoxystrobin) in spring and winter rapeseed production in Latvia [34] was the source of the largest contribution of 12% and thiophanate-methyl. In contrast, prochloraz (also systemic fungicide) in rapeseed production in Chile [52] was the source of the lowest contribution of 1%. Fungicides' contribution to freshwater ecotoxicity was higher than other environmental impacts irrespective of the dosage because the latter was the same order of magnitude in the cases of fruits and vegetables, and several authors grouped fungicides with pesticides during life cycle interpretation. A reason for the lower contributions of fungicides in cereals and rapeseed production can be the larger contribution of field emissions of chemical fertilizers because chemical fertilizers contribute to climate change, human toxicity, and ecotoxicity [35].

Finally, no distinction can be made regarding the impact of systemic and contact fungicides based on the reviewed studies. However, it should be pointed out that cereals and rapeseeds consisted of the only LCA studies that employed tebuconazole, epoxiconazole, and prochloraz, which are more toxic to mitochondrial activities, membrane degradations, and caspases 3/7 activities than insecticides and herbicides, according to [12].

3.4. Endpoint Impacts Due to Fungicides Application

In 2019, approximately 42,000, 55,430, and 255,870 tonnes of fruits, vegetables [53], and cereals [54], respectively, were produced in Europe with conventional farming [55,56]. It should be mentioned that non-edible parts of these agricultural products and rapeseed can be used for the production of biofuels [57,58] depending on the availability of money and consumers' habits [59]. Tables 2 and 3 show minimum and maximum benefits in human health per person and ecosystems quality (terrestrial), respectively, if fungicides are eliminated in fruits, vegetables, and cereals and rapeseed production in Europe. Outliers of climate change (in Figures 3 and 4) were not considered in endpoint calculations based on Grubbs' test. These outliers were derived from studies that grouped fungicides emissions with farm emissions.

Table 2. Human health and ecosystem quality (terrestrial) based on low contribution of fungicides to climate change, human toxicity, terrestrial ecotoxicity, and terrestrial acidification.

Disability Adjusted Life Years per Person (y·p ⁻¹)		Time Integrated Species Loss (Species·yr)		
Climate Change	Human Toxicity	Climate Change	Terrestrial Ecotoxicity	Terrestrial Acidification
Individualist perspective				
Fruits	4.4×10^{-7}	1.8×10^{-6}	0	0
Vegetables	4.6×10^{-7}	2.5×10^{-7}	0	7.7×10^{-1}
Cereals and rapeseed	1.8×10^{-7}	1.3×10^{-5}	0	6.2×10^{-1}
Hierarchical perspective				
Fruits	0	1.8×10^{-6}	0	0
Vegetables	0	2.5×10^{-7}	0	7.7×10^{-1}
Cereals and rapeseed	0	1.3×10^{-5}	0	6.2×10^{-1}
Egalitarian perspective				
Fruits	0	1.8×10^{-6}	0	0
Vegetables	0	2.5×10^{-7}	0	7.7×10^{-1}
Cereals and rapeseed	0	1.3×10^{-5}	0	6.2×10^{-1}

Table 3. Human health and ecosystem quality (terrestrial) based on high contribution of fungicides to climate change, human toxicity, terrestrial ecotoxicity, and terrestrial acidification.

Disability Adjusted Life Years per Person (y·p ⁻¹)		Time Integrated Species Loss (Species·yr)		
Climate Change	Human Toxicity	Climate Change	Terrestrial Ecotoxicity	Terrestrial Acidification
Individualist perspective				
Fruits	4.4×10^{-7}	1.8×10^{-5}	1.5	18.4
Vegetables	4.6×10^{-7}	2.7×10^{-6}	1.6	7.7
Cereals and rapeseed	1.8×10^{-7}	5.3×10^{-6}	0.6	3.1
Hierarchical perspective				
Fruits	5.0×10^{-6}	1.8×10^{-5}	7.7	18.4
Vegetables	5.3×10^{-6}	2.7×10^{-6}	8.2	7.7
Cereals and rapeseed	2.0×10^{-6}	5.3×10^{-6}	3.1	3.1
Egalitarian perspective				
Fruits	7.0×10^{-5}	1.8×10^{-5}	69.0	18.4
Vegetables	7.4×10^{-5}	2.7×10^{-6}	73.1	7.7
Cereals and rapeseed	2.8×10^{-5}	5.3×10^{-6}	28.1	3.1

Potential benefits for human health are greater in fruits than vegetables, cereals, and rapeseed. In addition, the fact that midpoint to endpoint conversion factors ranges minimally between individualist and hierarchist perspectives results in similar values for both perspectives for both low and high fungicides contributions to climate change and human toxicity. In contrast, concerning the egalitarian perspective, the climate change midpoint to endpoint conversion factor increases by at least one order of magnitude compared to the individualist perspective and results in adverse effects for all crops (Table 3). This increase in climate change conversion factor results in vegetables damaging human health more than cereals and rapeseed with the egalitarian perspective. Human toxicity is the midpoint indicator that influences human health results for all perspectives. Climate change insignificantly affects human health from individualist and hierarchist perspectives but increases its effect with the egalitarian perspective due to the increase

in midpoint to endpoint contribution. Lastly, the production of cereals is an order of magnitude higher than fruits and vegetables, but the very low contribution of fungicides in human toxicity of cereals and rapeseed does not result in higher toxicity per person than fruits.

The elimination of fungicides in cereals and rapeseed results in greater potential benefits in ecosystem quality (terrestrial) than vegetables and fruits for individualist and hierarchist perspectives. This occurs mainly due to production yields of cereals and rapeseed and larger terrestrial acidification score of cereals and rapeseed than other crops, especially with respect to the lowest amount of expected ecosystem benefits (Table 2). In contrast, terrestrial ecotoxicity is the largest contributor to benefits in ecosystem quality for the cases of fruits and vegetables based on the individualist and hierarchist perspectives. The individualist and hierarchist perspectives result in a greater difference in benefits among considered crops, but similar benefits are found with the egalitarian perspective. The egalitarian perspective results in more species loss due to the increase in midpoint to endpoint conversion factor for climate change and production yields for all crops.

Lastly, it should be mentioned that calculations for ecosystem quality (terrestrial) result in unrealistically high values of species loss. According to the International Union for Conservation of Nature, as of 2015, 36 species became extinct in Europe [60]. Thus, the term “species” here seems to be inappropriate, but “animals” could be used because the U.S. Fish and Wildlife Service estimates that approximately 67 million birds die from pesticide poisoning each year [61].

4. Conclusions

This review paper aimed to collect LCA studies focusing on the use of fungicides in fruits, vegetables, and cereals and rapeseed, and calculate damages to human health and ecosystem quality (terrestrial). It was found that LCA practitioners often do not distinguish the contribution of insecticides, herbicides, and fungicides to considered environmental impacts. Several LCA studies mentioned explicitly what fungicides are used and what is their dosage in the life cycle inventory stage, but grouped them with other pesticides in the life cycle interpretation stage.

It was found that fungicides contribute more to environmental impacts in the cultivation of fruits and vegetables than in the cultivation of cereals and rapeseed. Furthermore, fungicides dosage does not vary among the produced crops. Therefore, the specific chemical is crucial to be mentioned to select characterization factors to assess toxicity. If current conventional agriculture practices become organic, European citizens will benefit with up to 47 min per year of increased life quality due to fungicides elimination in fruits and vegetables cultivation. In addition, species loss benefits are also expected up to 90 species per year due to fungicides elimination in fruits and vegetables cultivation.

It is recommended that LCA practitioners increase transparency in their studies by showing what input materials are contributing to selected environmental impacts, and that new LCA studies are designed to assess explicitly the effect of different fungicides on human toxicity and ecotoxicity.

Supplementary Materials: The following supporting information can be downloaded at: <https://www.mdpi.com/article/10.3390/su14020846/s1>, Excel file.

Funding: We acknowledge support of this work by the project “AGRO4+ - Holistic approach to Agriculture 4.0 for new farmers” (MIS 5046239) which is implemented under the Action “Reinforcement of the Research and Innovation Infrastructure”, funded by the Operational Programme “Competitiveness, Entrepreneurship and Innovation” (NSRF 2014-2020) and co-financed by Greece and the European Union (European Regional Development Fund). The opinions expressed in this document reflect only the author’s view.

Institutional Review Board Statement: Not applicable.

Informed Consent Statement: Not applicable.

Data Availability Statement: The data presented in this study are available in Tables 2 and 3 within this article, and Supplementary Materials.

Acknowledgments: The author is grateful for the many constructive comments provided by the reviewers of earlier versions of this article.

Conflicts of Interest: The author declares no conflict of interest.

References

1. Ehrlich, P.R. A personal view: Environmental education—Its content and delivery. *J. Environ. Stud. Sci.* **2011**, *1*, 6–13. [\[CrossRef\]](#)
2. Muñoz, P.; Antón, A.; Núñez, M.; Paranjpe, A.; Ariño, J.; Castells, X.; Montero, J.; Rieradevall, J. Comparing The Environmental Impacts Of Greenhouse Versus Open-Field Tomato Production In The Mediterranean Region. *Acta Hortic.* **2008**, *801*, 1591–1596. [\[CrossRef\]](#)
3. Winans, K.; Brodt, S.; Kendall, A. Life cycle assessment of California processing tomato: An evaluation of the effects of evolving practices and technologies over a 10-year (2005–2015) timeframe. *Int. J. Life Cycle Assess.* **2019**, *25*, 538–547. [\[CrossRef\]](#)
4. D’Amico, G.; Szopik-Depczyńska, K.; Beltramo, R.; D’Adamo, I.; Ioppolo, G. Smart and Sustainable Bioeconomy Platform: A New Approach towards Sustainability. *Sustainability* **2022**, *14*, 466. [\[CrossRef\]](#)
5. D’Adamo, I.; Gastaldi, M.; Morone, P.; Rosa, P.; Sasanelli, C.; Settembre-Blundo, D.; Shen, Y. Bioeconomy of Sustainability: Drivers, Opportunities and Policy Implications. *Sustainability* **2021**, *14*, 200. [\[CrossRef\]](#)
6. OECD. *Food and Agriculture Organization of the United Nations OECD-FAO Agricultural Outlook 2021–2030*; OECD-FAO Agricultural Outlook; OECD: Paris, France, 2021; ISBN 978-92-64-43607-7.
7. Aktar, W.; Sengupta, D.; Chowdhury, A. Impact of pesticides use in agriculture: Their benefits and hazards. *Interdiscip. Toxicol.* **2009**, *2*, 1–12. [\[CrossRef\]](#) [\[PubMed\]](#)
8. Mazis, A.; Litskas, V.D.; Platis, D.P.; Menexes, G.C.; Anagnostopoulos, C.D.; Tsaboula, A.D.; Mamolos, A.P.; Kalburjji, K.L. Could energy equilibrium and greenhouse gas emissions in agroecosystems play a key role in crop replacement? A case study in orange and kiwi orchards. *Environ. Sci. Pollut. Res.* **2021**, *28*, 29421–29431. [\[CrossRef\]](#)
9. Esteve-Turrillas, F.; de la Guardia, M. Environmental impact of Recover cotton in textile industry. *Resour. Conserv. Recycl.* **2017**, *116*, 107–115. [\[CrossRef\]](#)
10. World Health Organization Pesticide Residues in Food. Available online: <https://www.who.int/news-room/fact-sheets/detail/pesticide-residues-in-food> (accessed on 18 December 2021).
11. Juraske, R.; Sanjuán, N. Life cycle toxicity assessment of pesticides used in integrated and organic production of oranges in the Comunidad Valenciana, Spain. *Chemosphere* **2011**, *82*, 956–962. [\[CrossRef\]](#)
12. Mesnage, R.; Defarge, N.; De Vendômois, J.S.; Séralini, G.-E. Major Pesticides Are More Toxic to Human Cells Than Their Declared Active Principles. *BioMed Res. Int.* **2014**, *2014*, 179691. [\[CrossRef\]](#)
13. Moharram, A.M.; Abdel-Hafez, S.I.I.; El-Said, A.H.M.; Salee, A. Effect of Two Systemic Fungicides on Cellulose Decomposing Fungi of Tomato Plants and on Some Enzymatic Activities. *Acta Microbiol. Immunol. Hung.* **2004**, *51*, 403–430. [\[CrossRef\]](#) [\[PubMed\]](#)
14. Sari, A.L.; Lubis, L. The effectiveness of contact fungicides mancozeb in controlling potato leaf blight disease (*Phytophthora infestans* (Mont) de Barry) in Karo District in the wet month and in the laboratory. In *IOP Conference Series: Earth and Environmental Science*; IOP Publishing: Bristol, UK, 2021; Volume 782.
15. Benbrook, C.; Kegley, S.; Baker, B. Organic Farming Lessens Reliance on Pesticides and Promotes Public Health by Lowering Dietary Risks. *Agronomy* **2021**, *11*, 1266. [\[CrossRef\]](#)
16. Gentil-Sargent, C.; Basset-Mens, C.; Renaud-Gentie, C.; Mottes, C.; Melero, C.; Launay, A.; Fantke, P. Introducing ground cover management in pesticide emission modeling. *Integr. Environ. Assess. Manag.* **2021**, *18*, 274–288. [\[CrossRef\]](#)
17. van Calker, K.J.; Berentsen, P.B.M.; De Boer, I.M.J.; Giesen, G.W.J.; Huirne, R.B.M. An LP-model to analyse economic and ecological sustainability on Dutch dairy farms: Model presentation and application for experimental farm “de Marke”. *Agric. Syst.* **2004**, *82*, 139–160. [\[CrossRef\]](#)
18. Wang, F.; Liu, Y.; Ouyang, X.; Hao, J.; Yang, X. Comparative environmental impact assessments of green food certified cucumber and conventional cucumber cultivation in China. *Renew. Agric. Food Syst.* **2018**, *33*, 432–442. [\[CrossRef\]](#)
19. He, X.; Qiao, Y.; Liu, Y.; Dendler, L.; Yin, C.; Martin, F. Environmental impact assessment of organic and conventional tomato production in urban greenhouses of Beijing city, China. *J. Clean. Prod.* **2016**, *134*, 251–258. [\[CrossRef\]](#)
20. Casolani, N.; Nissi, E.; Giampaolo, A.; Liberatore, L. Evaluating the effects of European support measures for Italian organic farms. *Land Use Policy* **2021**, *102*, 105225. [\[CrossRef\]](#)
21. Purnhagen, K.P.; Clemens, S.; Eriksson, D.; Fresco, L.O.; Tosun, J.; Qaim, M.; Visser, R.G.; Weber, A.P.; Wesseler, J.H.; Zilberman, D. Europe’s Farm to Fork Strategy and Its Commitment to Biotechnology and Organic Farming: Conflicting or Complementary Goals? *Trends Plant Sci.* **2021**, *26*, 600–606. [\[CrossRef\]](#)
22. ISO DIN EN ISO 14040:2006. *Environmental Management—Life Cycle Assessment—Principles and Framework*; ISO: Geneva, Switzerland, 2006.
23. ISO DIN EN ISO 14044:2006. *Environmental Management—Life Cycle Assessment—Requirements and Guidelines*, 1st ed.; ISO: Geneva, Switzerland, 2006.

24. Huijbregts, M.A.J.; Steinmann, Z.J.N.; Elshout, P.M.F.; Stam, G.; Verones, F.; Vieira, M.; Zijp, M.; Hollander, A.; van Zelm, R. ReCiPe2016: A harmonised life cycle impact assessment method at midpoint and endpoint level. *Int. J. Life Cycle Assess.* **2017**, *22*, 138–147. [\[CrossRef\]](#)

25. Van Zelm, R.; Huijbregts, M.; Posthuma, L.; Wintersen, A.; Van De Meent, D. Pesticide ecotoxicological effect factors and their uncertainties for freshwater ecosystems. *Int. J. Life Cycle Assess.* **2009**, *14*, 43–51. [\[CrossRef\]](#)

26. Dekker, E.; Zijp, M.C.; Van De Kamp, M.E.; Temme, E.H.M.; Van Zelm, R. A taste of the new ReCiPe for life cycle assessment: Consequences of the updated impact assessment method on food product LCAs. *Int. J. Life Cycle Assess.* **2019**, *25*, 2315–2324. [\[CrossRef\]](#)

27. Chatzisymeon, E.; Foteinis, S.; Borthwick, A.G.L. Life cycle assessment of the environmental performance of conventional and organic methods of open field pepper cultivation system. *Int. J. Life Cycle Assess.* **2017**, *22*, 896–908. [\[CrossRef\]](#)

28. Lopes, J.; Medeiros, D.L.; Kiperstok, A. Combining cleaner production and life cycle assessment for reducing the environmental impacts of irrigated carrot production in Brazilian semi-arid region. *J. Clean. Prod.* **2018**, *170*, 924–939. [\[CrossRef\]](#)

29. Romero-Gámez, M.; Antón, A.; Leyva, R.; Suárez-Rey, E.M. Inclusion of uncertainty in the LCA comparison of different cherry tomato production scenarios. *Int. J. Life Cycle Assess.* **2016**, *22*, 798–811. [\[CrossRef\]](#)

30. Pedretti, E.F.; Boakye-Yiadom, K.A.; Valentini, E.; Ilari, A.; Duca, D. Life Cycle Assessment of Spinach Produced in Central and Southern Italy. *Sustainability* **2021**, *13*, 10001. [\[CrossRef\]](#)

31. Tasca, A.L.; Nessi, S.; Rigamonti, L. Environmental Sustainability of Agri-Food Supply Chains: An LCA Comparison between Two Alternative Forms of Production and Distribution of Endive in Northern Italy. *J. Clean. Prod.* **2017**, *140*, 725–741. [\[CrossRef\]](#)

32. Russo, V.; Strever, A.E.; Ponstein, H.J. Exploring Sustainability Potentials in Vineyards through LCA? Evidence from Farming Practices in South Africa. *Int. J. Life Cycle Assess.* **2021**, *26*, 1374–1390. [\[CrossRef\]](#)

33. Abdul Rahman, M.H.; Chen, S.S.; Abdul Razak, P.R.; Abu Bakar, N.A.; Shahrun, M.S.; Zin Zawawi, N.; Muhamad Mujab, A.A.; Abdullah, F.; Jumat, F.; Kamaruzaman, R.; et al. Life Cycle Assessment in Conventional Rice Farming System: Estimation of Greenhouse Gas Emissions Using Cradle-to-Gate Approach. *J. Clean. Prod.* **2019**, *212*, 1526–1535. [\[CrossRef\]](#)

34. Fridrihsone, A.; Romagnoli, F.; Cabulis, U. Environmental Life Cycle Assessment of Rapeseed and Rapeseed Oil Produced in Northern Europe: A Latvian Case Study. *Sustainability* **2020**, *12*, 5699. [\[CrossRef\]](#)

35. Motevali, A.; Hashemi, S.J.; Tabatabaeekoloor, R. Environmental Footprint Study of White Rice Production Chain-Case Study: Northern of Iran. *J. Environ. Manag.* **2019**, *241*, 305–318. [\[CrossRef\]](#) [\[PubMed\]](#)

36. Câmara-Salim, I.; Almeida-García, F.; Feijoo, G.; Moreira, M.T.; González-García, S. Environmental Consequences of Wheat-Based Crop Rotation in Potato Farming Systems in Galicia, Spain. *J. Environ. Manag.* **2021**, *287*, 112351. [\[CrossRef\]](#) [\[PubMed\]](#)

37. Santos, I.V.; Bulle, C.; Levasseur, A.; Deschênes, L. Regionalized Terrestrial Ecotoxicity Assessment of Copper-Based Fungicides Applied in Viticulture. *Sustainability* **2018**, *10*, 2522. [\[CrossRef\]](#)

38. Khoshnevisan, B.; Rafiee, S.; Omid, M.; Mousazadeh, H.; Clark, S. Environmental Impact Assessment of Tomato and Cucumber Cultivation in Greenhouses Using Life Cycle Assessment and Adaptive Neuro-Fuzzy Inference System. *J. Clean. Prod.* **2014**, *73*, 183–192. [\[CrossRef\]](#)

39. Abeliotis, K.; Detsis, V.; Pappia, C. Life Cycle Assessment of Bean Production in the Prespa National Park, Greece. *J. Clean. Prod.* **2013**, *41*, 89–96. [\[CrossRef\]](#)

40. Avadí, A.; Marcin, M.; Biard, Y.; Renou, A.; Gourlot, J.-P.; Basset-Mens, C. Life Cycle Assessment of Organic and Conventional Non-Bt Cotton Products from Mali. *Int. J. Life Cycle Assess.* **2020**, *25*, 678–697. [\[CrossRef\]](#)

41. Shen, X.; Zhang, L.; Zhang, J. Ratoon Rice Production in Central China: Environmental Sustainability and Food Production. *Sci. Total Environ.* **2021**, *764*, 142850. [\[CrossRef\]](#)

42. Roy, P.-O.; Azevedo, L.B.; Margni, M.; van Zelm, R.; Deschênes, L.; Huijbregts, M.A.J. Characterization Factors for Terrestrial Acidification at the Global Scale: A Systematic Analysis of Spatial Variability and Uncertainty. *Sci. Total Environ.* **2014**, *500–501*, 270–276. [\[CrossRef\]](#)

43. Lu, T.; Zhang, Q.; Lavoie, M.; Zhu, Y.; Ye, Y.; Yang, J.; Paerl, H.W.; Qian, H.; Zhu, Y.-G. The Fungicide Azoxystrobin Promotes Freshwater Cyanobacterial Dominance through Altering Competition. *Microbiome* **2019**, *7*, 128. [\[CrossRef\]](#)

44. Gaspar, P.D.; Godina, R.; Barrau, R. Influence of Orchard Cultural Practices during the Productive Process of Cherries through Life Cycle Assessment. *Processes* **2021**, *9*, 1065. [\[CrossRef\]](#)

45. Beauchet, S.; Rouault, A.; Thiollet-Scholtus, M.; Renouf, M.; Jourjon, F.; Renaud-Gentie, C. Inter-Annual Variability in the Environmental Performance of Viticulture Technical Management Routes—a Case Study in the Middle Loire Valley (France). *Int. J. Life Cycle Assess.* **2019**, *24*, 253–265. [\[CrossRef\]](#)

46. Mohseni, P.; Borghei, A.M.; Khanali, M. Coupled Life Cycle Assessment and Data Envelopment Analysis for Mitigation of Environmental Impacts and Enhancement of Energy Efficiency in Grape Production. *J. Clean. Prod.* **2018**, *197*, 937–947. [\[CrossRef\]](#)

47. Ilari, A.; Toscano, G.; Boakye-Yiadom, K.A.; Duca, D.; Foppa Pedretti, E. Life Cycle Assessment of Protected Strawberry Productions in Central Italy. *Sustainability* **2021**, *13*, 4879. [\[CrossRef\]](#)

48. Naderi, S.; Ghasemi Nejad Raini, M.; Taki, M. Measuring the Energy and Environmental Indices for Apple (Production and Storage) by Life Cycle Assessment (Case Study: Semiroom County, Isfahan, Iran). *Environ. Sustain. Indic.* **2020**, *6*, 100034. [\[CrossRef\]](#)

49. Guo, X.-X.; Zhao, D.; Zhuang, M.-H.; Wang, C.; Zhang, F.-S. Fertilizer and Pesticide Reduction in Cherry Tomato Production to Achieve Multiple Environmental Benefits in Guangxi, China. *Sci. Total Environ.* **2021**, *793*, 148527. [\[CrossRef\]](#)

50. Gil, R.; Bojacá, C.R.; Schrevens, E. Does Optimized Agrochemicals Management Help to Reduce the Environmental Impact in Tomato Production? A Comparative Analysis between Greenhouse and Open Field Systems. In Proceedings of the Acta Horticulturae: 2020, Leuven, Belgium, 23 November 2020; International Society for Horticultural Science (ISHS): Leuven, Belgium, 2020; pp. 1145–1152.

51. Yelboğa, M.N.M. LCA Analysis of Grafted Tomato Seedling Production in Turkey. *Sustainability* **2020**, *12*, 25. [CrossRef]

52. Iriarte, A.; Rieradevall, J.; Gabarrell, X. Environmental Impacts and Energy Demand of Rapeseed as an Energy Crop in Chile under Different Fertilization and Tillage Practices. *Biomass Bioenergy* **2011**, *35*, 4305–4315. [CrossRef]

53. Messe Berlin GmbH. *Fruitnet Fruit Logistica*. In *European Statistics Handbook*; Messe Berlin GmbH: Berlin, Germany, 2021.

54. Eurostat Agricultural Production—Crops. Available online: https://ec.europa.eu/eurostat/statistics-explained/index.php?title=Agricultural_production_-_crops (accessed on 17 December 2021).

55. Röös, E.; Mie, A.; Wivstad, M.; Salomon, E.; Johansson, B.; Gunnarsson, S.; Wallenbeck, A.; Hoffmann, R.; Nilsson, U.; Sundberg, C.; et al. Risks and Opportunities of Increasing Yields in Organic Farming: A Review. *Agron. Sustain. Dev.* **2018**, *38*, 14. [CrossRef]

56. Trávníček, J.; Schaack, D.; Willer, H. Organic Agriculture in Europe: Current Statistics; 2021. Available online: https://www.google.com/url?sa=t&rct=j&q=&esrc=s&source=web&cd=&ved=2ahUKEwisloXJ6qv1AhWCy6QKHT9ID7cQFnoECBYQAQ&url=https%3A%2F%2Forgprints.org%2F39367%2F1%2Ftravnicek-etal-2021-02-18-2pm-european-market-final.pdf&usg=AOvVaw00LPk_mELOPe2K_2eMZ5tA (accessed on 17 December 2021).

57. Ikram, M.; Sroufe, R.; Awan, U.; Abid, N. Enabling Progress in Developing Economies: A Novel Hybrid Decision-Making Model for Green Technology Planning. *Sustainability* **2022**, *14*, 258. [CrossRef]

58. Ding, Z.; Grundmann, P. Development of Biorefineries in the Bioeconomy: A Fuzzy-Set Qualitative Comparative Analysis among European Countries. *Sustainability* **2022**, *14*, 90. [CrossRef]

59. Bekti, D.B.M.; Prasetyo, Y.T.; Redi, A.A.N.P.; Budiman, A.S.; Mandala, I.M.P.L.; Putra, A.R.; Persada, S.F.; Nadlifatin, R.; Young, M.N. Determining Factors Affecting Customer Intention to Use Rooftop Solar Photovoltaics in Indonesia. *Sustainability* **2022**, *14*, 280. [CrossRef]

60. International Union for Conservation of Nature. European Species Under Threat. In *Overview of European Red Lists Results*; International Union for Conservation of Nature: Gland, Switzerland, 2015.

61. Mineau, P. Direct Losses of Birds to Pesticides—Beginnings of a Quantification. In *USDA Forest Service General Technical Report*; Taylor & Francis: Abingdon, UK, 2005.