

**Natronosporangium hydrolyticum gen. nov., sp. nov., a haloalkaliphilic polyhydrolytic actinobacterium from a soda solonchak soil in Central Asia**

Sorokin, Dmitry Y.; Elcheninov, Alexander G.; Khijniak, Tatiana V.; Zaharycheva, Alicia P.; Boueva, Olga V.; Ariskina, Elena V.; Bunk, Boyke; Spröer, Cathrin; Evtushenko, Lyudmila I.; More Authors

**DOI**

[10.1016/j.syapm.2022.126307](https://doi.org/10.1016/j.syapm.2022.126307)

**Publication date**

2022

**Document Version**

Final published version

**Published in**

Systematic and Applied Microbiology

**Citation (APA)**

Sorokin, D. Y., Elcheninov, A. G., Khijniak, T. V., Zaharycheva, A. P., Boueva, O. V., Ariskina, E. V., Bunk, B., Spröer, C., Evtushenko, L. I., & More Authors (2022). *Natronosporangium hydrolyticum* gen. nov., sp. nov., a haloalkaliphilic polyhydrolytic actinobacterium from a soda solonchak soil in Central Asia. *Systematic and Applied Microbiology*, 45(3), Article 126307. <https://doi.org/10.1016/j.syapm.2022.126307>

**Important note**

To cite this publication, please use the final published version (if applicable).  
Please check the document version above.

**Copyright**

Other than for strictly personal use, it is not permitted to download, forward or distribute the text or part of it, without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license such as Creative Commons.

**Takedown policy**

Please contact us and provide details if you believe this document breaches copyrights.  
We will remove access to the work immediately and investigate your claim.



## *Natronosporangium hydrolyticum* gen. nov., sp. nov., a haloalkaliphilic polyhydrolytic actinobacterium from a soda solonchak soil in Central Asia

Dimitry Y. Sorokin<sup>a,b,\*</sup>, Alexander G. Elcheninov<sup>a</sup>, Tatiana V. Khijniak<sup>a</sup>, Alicia P. Zaharycheva<sup>a</sup>, Olga V. Boueva<sup>c</sup>, Elena V. Ariskina<sup>c</sup>, Boyke Bunk<sup>d</sup>, Cathrin Spröer<sup>d</sup>, Lyudmila I. Evtushenko<sup>c</sup>, Ilya V. Kublanov<sup>a</sup>, Richard L. Hahnke<sup>d</sup>

<sup>a</sup> Winogradsky Institute of Microbiology, Research Centre of Biotechnology, Russian Academy of Sciences, Moscow, Russia

<sup>b</sup> Department of Biotechnology, TU Delft, the Netherlands

<sup>c</sup> Leibniz Institute DSMZ – German Collection of Microorganisms and Cell Cultures, Germany

<sup>d</sup> Institute of Biochemistry and Physiology of Microorganisms, Russian Academy of Sciences, Puschino, Moscow Region, Russia

### ARTICLE INFO

#### Article history:

Received 21 September 2021

Revised 1 February 2022

Accepted 7 February 2022

#### Keywords:

Soda solonchak soils  
Haloalkaliphilic  
Actinobacteria  
*Micromonosporaceae*  
Polyhydrolytic

### ABSTRACT

During a cultural diversity survey on hydrolytic bacteria in saline alkaline soils, a hydrolytic actinobacterium strain ACPA39<sup>T</sup> was enriched and isolated in pure culture from a soda solonchak soil in southwestern Siberia. It forms a substrate mycelium with rod-shaped sporangia containing 1–3 exospores. The isolate is obligately alkaliphilic, growing at pH 7.5–10.3 (optimum at 8.5–9.0) and moderately halophilic, tolerating up to 3 M total Na<sup>+</sup> in the form of sodium carbonates. It is an obligately aerobic, organoheterotrophic, saccharolytic bacterium, utilizing various sugars and alpha/beta-glucans as growth substrates. According to the 16S rRNA gene-based phylogenetic analysis, strain ACPA39<sup>T</sup> forms a distinct branch within the family *Micromonosporaceae*, with the sequence identities below 94.5% with type strains of other genera. This is confirmed by phylogenomic analysis based on the 120 conserved single copy protein-based markers and genomic indexes (ANI, AAI). The cell-wall of ACPA39<sup>T</sup> contained meso-DAP, glycine, glutamic acid and alanine in a equimolar ratio, characteristic of the peptidoglycan type A1γ'. The whole-cell sugars include galactose and xylose. The major menaquinone is MK-10(H<sub>4</sub>). The identified polar lipids consist of phosphatidylglycerol, diphosphatidylglycerol, phosphatidylethanolamine and phosphatidylinositol. The polar lipid fatty acids were dominated by *anteiso*-C<sub>17:0</sub>, *iso*-C<sub>16:0</sub>, *iso*-C<sub>17:0</sub>, 10 Me-C<sub>18:0</sub> and C<sub>18:1</sub>ω9. Based on the distinct phylogeny, the chemotaxonomy features and unique phenotypic properties, strain ACPA39<sup>T</sup> (DSM 106523<sup>T</sup> = VKM 2772<sup>T</sup>) is classified into a new genus and species in the family *Micromonosporaceae* for which the name *Natronosporangium hydrolyticum* gen. nov., sp. nov. is proposed.

© 2022 The Author(s). Published by Elsevier GmbH. This is an open access article under the CC BY license (<http://creativecommons.org/licenses/by/4.0/>).

### Introduction

Actinobacteria represent one of the dominant and well characterized phylum in soils. In contrast to eukaryotic fungi, many of the actinobacterial members are rather alkalitolerant, with that trait being especially common among the numerous *Streptomyces* and *Nocardiopsis* species [1–3]. Saline alkaline soils (soda solonchaks) with stable high pH above 9 [4–6] are ideal habitats for such actinobacteria due to a domination of sodium carbonates among the soluble salts but, currently, the prokaryotic

communities of this extreme habitat are poorly investigated. To date, only a few publications report on the abundance of alkali-tolerant actinomycetes belonging to genera *Streptomyces* and *Nocardiopsis* [7–12].

Our recent investigation of the diversity of culturable bacteria with hydrolytic potential inhabiting soda solonchak soils from different geographical locations in Central Asia yielded a large proportion of haloalkaliphilic actinobacteria, mainly belonging to the mycelium-forming genera *Streptomyces* and *Nocardiopsis*, and a less abundant single-cell isolates belonging to the genus *Isoptericola* [13]. In addition, two isolates clearly differed from the known haloalkaliphilic species forming newgenera in the families *Glycomycetaceae* and *Micromonosporaceae*. The former has recently

\* Corresponding author.

E-mail addresses: [soroc@inmi.ru](mailto:soroc@inmi.ru), [d.sorokin@tudelft.nl](mailto:d.sorokin@tudelft.nl) (D.Y. Sorokin).

been described as *Natronoglycomyces albus* [14]. The purpose of this work was to classify the novel haloalkaliphilic member of the *Micromonosporaceae* family, strain ACPA39<sup>T</sup>.

## Materials and methods

### Enrichment and cultivation conditions

Strain ACPA39<sup>T</sup> was isolated from surface horizon (0–5 cm deep) of a soda solonchak soil in southwestern Siberia (Barabinskaya Steppe, Novosibirsk region, Russia; 55.06 N/81.1 E). The 1:5 water extract had a pH of 10.7, a total soluble salts content of 62 g kg<sup>-1</sup> and a total soluble carbonate alkalinity of 0.51 M. Enrichment and isolation of a pure culture was performed in a sodium carbonate medium containing 0.6 M total Na<sup>+</sup> at pH 10 with carboxymethylcellulose (1 g l<sup>-1</sup>) and yeast extract (0.2 g l<sup>-1</sup>) as substrates, as described previously [13,14]. Growth and cell/colony morphology were investigated in cultures grown during 2–28 days at 28 °C in liquid and solid media including: 1) soluble starch/yeast extract medium (SS-YE) (1 g l<sup>-1</sup> starch/0.2 g l<sup>-1</sup> yeast extract; pH 10 and 0.6 M total Na<sup>+</sup>); 2) agar media mixed (1:1 v/v) with sodium carbonate base buffer (15 g Na<sub>2</sub>CO<sub>3</sub>, 20 g NaHCO<sub>3</sub>, 3 g NaCl, 1 g K<sub>2</sub>HPO<sub>4</sub> per liter, pH 9.5) and ISP 2 [15], R2A [16] or PYG medium (5 g peptone, 3 g yeast extract and 5 g glucose per liter). The growth pH profiling was performed with the following buffer systems (containing 0.6 M total Na<sup>+</sup>): 50 mM potassium phosphate-HEPES/0.6 M NaCl for pH 5–8, 50 mM HEPES-0.6 M NaHCO<sub>3</sub> for pH 8–8.5 and sodium carbonate-bicarbonate for pH 9–11 (the step was 0.5 pH unit). The final pH were used as the actual values [11]. It must be argued, however, whether the pH profiling in some literature was performed convincingly [10,12], since the recommended use of strong buffering systems in liquid media compatible with biology and continuous monitoring of the pH changes during growth was not applied [11]. Thus, reported pH maxima of 12 are unrealistic, because there are no buffering systems maintaining stable pH above 11.

The influence of Na<sup>+</sup> (in the form of sodium carbonates, between 0 and 4 M, pH 10) and temperature (within the range from 10 to 60 °C with intervals of 5 °C; pH 10 and 0.6 M Na<sup>+</sup>) on growth were investigated with starch/yeast extract-containing medium. For the N-source spectrum, the culture was grown in liquid medium at pH 9.5/0.6 M Na<sup>+</sup> and 30 °C with addition of ammonium, nitrate or urea at 4 mM total N or 0.2 g l<sup>-1</sup> of yeast extract peptons from meat or casein. For the antimicrobial activity tests, paper discs were soaked in the filter-sterilized culture supernatant of strain ACPA39<sup>T</sup> pregrown for two weeks on starch/yeast extract medium and the discs were placed on the surface of the LB agar preinoculated with either *E. coli* or *Micrococcus luteus* test cultures.

### Cell morphology and chemotaxonomy

Cell micromorphology was studied with light microscopy (Zeiss Axioplan Imaging 2, Göttingen, Germany) and transmission and scanning electron microscopy (JEOL, Japan). For thin section electron microscopy, the cell material was fixed with paraformaldehyde (3% w/v in 0.6 M NaCl) for 2 h at room temperature, postfixed by OsO<sub>4</sub> (1% w/v in 0.6 M NaCl) for 48 h at 4 °C, embedded into agar blocks, stained with uranyl acetate (1% w/v) for 1 h, dehydrated in alcohol series and finally in absolute acetone, embedded into Epoxy resin and thin sectioned on the ultramicrotome. The ultrathin sections were finally stained with lead citrate (1% w/v) for 1 h.

For chemotaxonomic characterization cells were grown in sodium carbonate medium at pH 9.5 and 0.6 M total Na<sup>+</sup>, harvested

in late exponential growth phase and freeze-dried. The cell wall fraction was obtained by hydrolysis in 0.1 M NaOH [17] followed with hydrolysis with 6 M HCl at 110 °C for 20 h. The isomer composition of diaminopimelic acid (DAP) was determined by TLC. The cell wall amino acids composition was analyzed by anion-exchange chromatography (Biotronik, Germany) equipped with the DC6a resin column (Durrum) [18]. The whole cell sugar composition was determined by anion-exchange chromatography [Biotronik, Germany; column DA-x8-11 (Durrum)] after hydrolysis of cells in 3 M trifluoroacetic acid at 100 °C for 6 h [19]. Intact membrane polar lipids (MPL) were extracted by acidic methanol and separated by two-dimensional TLC using HPTLC Silica gel 60, chloroform-methanol-water (65:25:4) system in the first direction and chloroform-acetic acid-methanol-water (80:15:12:4) in the second direction [20]. The identified MPL spots were compared with respective spots on chromatograms of related reference strains for which the MPL profiles were determined previously. Polar lipid fatty acids (PLFA) were extracted, saponified, methylated and purified, and the methyl esters were analyzed by a Trace GC Ultra coupled to a DSQ II single-quadrupole mass spectrometer (Thermo Scientific) and identified by using the NIST 17 mass spectral library (<https://chemdata.nist.gov/dokuwiki/doku.php?id=c hemdata:nist17>) [21]. Respiratory menaquinones were extracted from wet cells, purified according to Collins (1985) [22] and analyzed with a LCQ Advantage MAX mass spectrometer (Thermo Finnigan).

### Hydrolytic activity

The potential of ACPA33<sup>T</sup> to hydrolyze various polysaccharides was investigated at three different levels: (1) by the ability to use the polysaccharides as growth substrates; (2) by the hydrolytic activity assays either on agar plates with zone hydrolysis detection [13] or by incubation of cell suspensions in titer plates with the dye conjugated polymers as described previously for *Natronoglycomyces albus* [14]; (3) by querying the genome for the presence of glycosyl hydrolase (GH)-encoding genes using the dbCAN2 server [23].

### Genomic sequencing, phylogenetic and genomic analyses

Genomic DNA was extracted according to Park [24]. SMRTbell<sup>TM</sup> template library was prepared using SMRTbell<sup>®</sup> Express Template Prep Kit 2.0 PacificBiosciences (CA, USA). For preparing 10 kb libraries, 1 µg genomic DNA was end-repaired and ligated to bar-coded adapters. The amplified products were pooled and sequenced in 1/16 SMRT cell on the Sequel<sup>III</sup> (PacificBiosciences, Menlo Park, CA, USA) resulting in 704,432 aligned subreads with mean aligned read lengths of 4.821 bp. Long read genome assembly was performed with the “Microbial Assembly” protocol included in SMRTlink version 8 using default parameters with exception of the target genome size, which was set to 5,7 Mbp. The assembly produced a single circular chromosome and two circular plasmids were obtained, afterwards rotated to the chromosomal/plasmid origin (*dnaA*, *parA*). Identification of replication genes has been done based on BLAST.

Primary identification of strain ACPA39<sup>T</sup> was performed using NCBI BLAST with one of two copies of the 16S rRNA gene sequence as the query.

For phylogenetic analysis based on comparison of 16S rRNA genes, the 16S rRNA gene sequence of strain ACPA39<sup>T</sup>, the sequences of all species with validly published names within the *Micromonosporaceae* family as well as *Rubrobacter radiotolerans* P1 as an outgroup were aligned using MAFFT server with G-INS-i method [25]. The phylogenetic tree was constructed in MEGA7 [26] using the maximum likelihood method and the

GTR model (G + I, 4 categories) with 1000 bootstrap replications; all positions with less than 95% site coverage were eliminated. For extended phylogenomic analysis based on the “bac120” set of conserved single copy bacterial proteins [27], the protein sequences were identified and aligned in *in silico* proteomes of type species of each genus within *Micromonosporaceae* (*Catenulanes* genus was excluded from analysis due to a bad quality of genome assemblies; for two multispecies genera, *Micromonospora* and *Actinoplanes*, several most distant species were selected using the GTDB-tk v.1.6.0 with reference data v.202) [28]. The alignment was treated using the trimAl v.1.4.1 with the -gt 1 parameter (full gap elimination) [29]. The phylogenomic tree was constructed in the RAxML v.8.2.12 [30] with the PROTGAM-MAILG model of amino acid substitution; local support values were 1000 rapid bootstrap replications. Phylogenetic trees were visualized using iTOL v.6.3.1 [31]. The whole genome-based comparisons were performed as follows: pairwise ANI values were calculated using the *pyani* module v.0.2.8 [32] with ANIb method; AAI values – using AAI matrix calculator [33] with Diamond program [34]. The Percentage of Conserved Proteins (POCP) values were calculated using the following script: (<https://github.com/hoelzer/pocp>).

## Results

### Morphological and chemotaxonomy features

During growth on solid medium at pH 9.5 with starch and yeast extract, strain ACPA39<sup>T</sup> formed mostly substrate mycelium varying in color from grayish to brown, depending on age. In old cultures, copious number of conidia were branching off the vegetative mycelium, first in a common sporangium (as was obvious from the thin-section electron microscopy), single or in rosette-like clusters, and in a later growth stage fragmenting into two to four angular sporangiospores with a slightly rough surface. The sporangia were also formed in liquid cultures (Fig. 1). No diffusible pigment was produced on any of the solid media tested.

The amino acids diagnostic for peptidoglycan detected in the cell wall of ACPA39<sup>T</sup> included diamino acid meso-DAP (peptidoglycan type A1γ') and also glycine, glutamate and alanine in approximately equal proportions (1 : 1 : 1 : 1 : 0.7). The cell-wall sugars included galactose and xylose. The identified MPL included diphosphatidylglycerol (DPG), phosphatidylethanolamine (PE), phosphatidylinositol (PI) and phosphatidylinositol mannoside (PIM). Those MPL species are also present in most of the *Micromonosporaceae* genera (Table 1). In addition, unidentified MPL were also present, including four phospho-, four glyco- and two aminoglycolipids (Supplementary Fig. S1). The dominant PLFA were anteiso-C<sub>17:0</sub> and iso-C<sub>16:0</sub>, iso-C<sub>17:0</sub>, 10 Me-C<sub>18:0</sub> and C<sub>18:1</sub>ω9 (45, 15, 7, 6.7 and 5.6% from the total, respectively). The dominant respiratory menaquinone was identified as MK-10(H<sub>4</sub>) (Supplementary Fig. S2), which is often present in various actinobacteria and members of the *Micromonosporaceae* in particular [35,36], while the absence of MK-9 species is not as common (Table 1).

### Genome sequencing

The genome of ACP39<sup>T</sup> was sequenced and assembled until completeness in the DSMZ. The final genome sequence is available in the GenBank under the numbers CP070499.1. Raw sequencing read data were deposited at NCBI SRA (698925). The genome size is 5.7 Mbp consisting of 5,177 genes encoding for 5,037 proteins. There are two *rrm* operons with identical sequences.

### Phylogenetic analysis

According to 16S rRNA gene sequence BLAST search, strain ACPA39<sup>T</sup> is a member of the *Actinobacteria*, family *Micromonosporaceae* [36,37] with the highest 16S rRNA gene sequence identity around 94% with the type strain CNH643<sup>T</sup> of marine species *Salinispora arenicola* and *Actinoplanes subglobosus*. 16S rRNA gene sequence-based phylogenetic analysis showed that ACPA39<sup>T</sup> forms a separate branch within the family *Micromonosporaceae* but several key nodes are not significantly supported. A search for more closely related 16S rRNA gene sequences among the uncultured database identified a few clones with sequence identities 97–98% to ACPA39<sup>T</sup>, indicating that they might belong to the same new genus as ACPA39<sup>T</sup>. In particular, two clones from the saline alkaline lakes Texcoco and Alchichica in Mexico (TX1A 103 and Alchichica AQ22 1B 75) were the closest to ACPA39<sup>T</sup> (Supplementary Fig. S3).


A more advanced phylogenomic approach based on 120 conserved protein markers (bac120 set) yielded a tree with a more stable branching order and revealed that ACPA39<sup>T</sup> forms a novel genus-level lineage clustering in an independent group with the genus *Pilimelia* as the closest relative. This branch is positioned between two large monophyletic clusters, one with the genus *Catellatospora*, together with its closely related genera and the other with the multispecies genus *Actinoplanes* (Fig. 2). A search of the metagenomic databases did not identify any apparent uncultured members belonging to the ACPA39<sup>T</sup> lineage, most probably because saline soda soils are not yet represented in such databases (Fig. 2).

The calculated AAI values between ACPA39<sup>T</sup> and members of the related genera from the *Micromonosporaceae* with available genome sequences ranged from 55.8% to 62.8% (Table S1a), which is below 65%, the threshold proposed for different genera [38]. The POCP values were 40.9–51.5% (Table S1b); thus maximum PCOP between the strain ACPA39<sup>T</sup> and relatives was slightly higher than the genus-level boundary – 50% [39]. But there are many works revealing that strict boundary of 50% was not appropriate for genera delineation within some taxa [40–45]. The ANI values (72.4–75.3%) were close to the median and mean intergeneric values (75.5% and 75.7%, respectively) calculated for representatives of this family (Table S1c).

Moreover, numerous probable misclassifications in the family that would need to be addressed in the future were uncovered using the phylogenomic approach. These include the genera *Couchioplanes*, *Krasilnikovia*, *Mangrovihabitans* and *Pseudosporangium*, which most probably belong to the genus *Actinoplanes*; the genera *Allorhizocola* and *Rhizocola* – as species of the genus *Catelliglobospora*; the genera *Plantactinospira*, *Polymorphospira* and *Salinispora* – as species of the type genus *Micromonospora*.

### Growth physiology and hydrolytic potential

Growth at different pH values and Na-carbonate concentrations was assessed on starch/yeast extract medium at 0.6 M total Na<sup>+</sup> (for variable pH) and at pH 10 (for different salinities). The strain grew optimally at low salt (0.1–0.3 M Na<sup>+</sup>) but with the high maximum tolerance of up to 3 M Na<sup>+</sup> (Fig. 3a). ACPA39<sup>T</sup> started to grow actively within a pH range from 7.5 to 10.3 (these are the final pH values) and had a moderately alkaline optimum at pH 8.2–9.0 (Fig. 3b), which qualifies this organism as an obligate alkaliphile. The genome search identified a gene locus coding for a single (multisubunit) sodium/proton antiporter MrpA-B/CDEFG as prerequisite for maintaining a near-neutral pH inside the cells of alkaliphiles [46]. *Salinispora arenicola* is the only member of the family *Micromonosporaceae* reported as able to grow at extremely high pH values up to 12, despite being isolated from a pH-neutral habitat [12]. However, as mentioned in the introduction,



**Fig. 1.** Cell morphology of strain ACPA39<sup>T</sup> growing on starch/yeast extract medium at pH 10 and 0.6 M total Na<sup>+</sup>. (a–b), phase contrast microscopy showing formation of sporangia in young cultures (a) and defragmentation of sporangia into separate spores in old culture (b); (c, g), scanning electron microphotographs showing sporangia and separate spores; (d–f, h), thin section electron microscopy showing a sporangium with spores (d), release of spores from sporangium (e) and separated mature exospores (h).

**Table 1**  
Phenotypic comparison of haloalkaliphilic strain ACPA39<sup>T</sup> from soda soils with the phylogenetically closely related genera in the family *Micromonosporaceae* (data from [12,36,47–53]).

Property	<b>"<i>Natronosporangium</i>"</b>	<i>Pilimelia</i>	<i>Catellatospora</i>	<i>Catelliglobospora</i>	<i>Hamadaea</i>	<i>Actinoplanes</i>	<i>Salinispora</i>
Number of species with validly published names	1	3	8	1	2	50	9
Aerial mycelium	Gray to brown, poor	Yellow, poor	No	No	No	No or rudimentary	No
Substrate mycelium	Abundant, branching, brown	Abundant, yellow-orange	Abundant, yellow	Abundant, yellow	Abundant, yellow	Abundant, color variable with orange as dominant	Abundant, branching, from orange to black
Exospores	2–4 in sporangia, angular, 0.2–0.6 µm, slightly rough	Sporangia with multiple spores, motile	Chains without sporangia	Chains without sporangia	Chains without sporangia	Multiple globular in sporangia, motile; 1.1–1.6 µm diameter	In grape-like bunches, round, 0.8–3.8 µm, smooth
Peptidoglycan diagnostic amino acids	<i>meso</i> -DAP: glutamate : glycine: alanine (1 : 1 : 1 : 0.7)	<i>meso</i> -DAP; glycine	<i>meso</i> -DAP; 3-OH- <i>meso</i> -DAP; glycine	<i>meso</i> -DAP; glycine	<i>meso</i> -DAP; 3-OH- <i>meso</i> -DAP; glycine	<i>meso</i> -DAP, glycine; 3-OH- <i>meso</i> -DAP (V)	<i>meso</i> -DAP; glycine
Cell wall sugars	Galactose, xylose	Arabinose, xylose	galactose, xylose, arabinose, ribose, glucose, mannose; rhamnose (V)	galactose, xylose, arabinose, glucose, mannose, rhamnose	galactose, xylose, arabinose, ribose, glucose, mannose; rhamnose	Xylose; galactose, arabinose (V)	galactose, xylose; arabinose, ribose, glucose (V)
Identified polar lipids	PG, DPG, PE, PI	PE, PI	DPG, PE, PI; PIM (V)		DPG, PE, PI, PIM; PME(V)	PE; DPG, PG, PI, PIM (V)	DPG, PE, PI; PME; PG and PIM (V)
Dominant menaquinones	10(H <sub>4</sub> )	9(H <sub>2</sub> ,H <sub>4</sub> )	10(H <sub>6</sub> ;H <sub>8</sub> ); 9(H <sub>4</sub> ; H <sub>6</sub> )	10(H <sub>4</sub> )	9(H <sub>6</sub> )	9(H <sub>4</sub> ); 9(H <sub>6</sub> ) (V)	9(H <sub>4</sub> )
Dominant PLFA (in order of abundance)	<i>anteiso</i> -C <sub>17:0</sub> , <i>iso</i> -C <sub>16:0</sub>	C <sub>15:0</sub> ; <i>iso</i> -C <sub>15:0</sub> ; C <sub>17:0</sub>	C <sub>17:0</sub> , C <sub>17:1</sub> Ω8c, <i>iso</i> -C <sub>15:0</sub> , <i>iso</i> -C <sub>16:0</sub>	C <sub>17:0</sub> , <i>iso</i> -C <sub>15:0</sub> , <i>iso</i> -C <sub>16:0</sub>	<i>iso</i> -C <sub>15:0</sub> , <i>anteiso</i> -C <sub>17:0</sub> ; <i>iso</i> -C <sub>16:0</sub> ; <i>anteiso</i> -C <sub>15:0</sub>	<i>iso</i> -C <sub>15:0</sub> , <i>iso</i> -C <sub>16:0</sub> , <i>anteiso</i> -C <sub>17:0</sub>	<i>iso</i> -C <sub>16:0</sub> , <i>iso</i> -C <sub>15:0</sub>
Polysaccharide as growth substrates	starch, dextrin, cyclodextrin, glycogen, pullulan, inulin, amorphous cellulose, laminarin, lichenan, barley glucan, xylane, pachyman, beta-mannan, glucomannan, xyloglycan, arabinoxylan	-	Starch (V)	Starch	Starch	Starch/dextrin, chitin, pectin (multiple); cellulose (few)	Starch, chitin
Sugar utilized for growth	Glucose, galactose, rhamnose, dextrose, arabinose, maltose, sucrose, cellobiose, trehalose, melizitose, raffinose, lactose	-	Glucose, galactose, arabinose; maltose, melibiose, ribose, rhamnose, lactose, mannose, sucrose, trehalose, xylose, mannitol (all V)	Glucose, galactose, arabinose; maltose, melibiose, ribose, rhamnose, lactose, mannose, sucrose, trehalose, xylose, mannitol (V)	Glucose, arabinose, melibiose, xylose, rhamnose, mannose, sucrose, trehalose; galactose, ribose, maltose, raffinose, fructose, cellobiose (V)	Glucose, galactose, fructose, arabinose, mannose, maltose, melibiose, xylose, sucrose, trehalose; lactose, melizitose, raffinose, rhamnose, ribose, cellobiose sorbitol, inositol, mannitol (V)	Cellobiose, melizitose, lactose
Proteolytic activity	Casein, gelatin	Keratine, casein, gelatin (V)	Gelatin (V)	Casein, gelatine	Gelatin (V)	Casein, gelatine	Casein, elastin, gelatin: (all V)
Lipase activity	- (olive oil)	NR	NR	NR	+Tween80	+Tween80 (NR in most species)	V (Tween80)
Oxidase/catalase	w/+	NR/NR	NR/+	+/+	+/+	V/V	V/+
pH range (optimum)	7.5–10.3 (8.5)	5.0–7.8 (6.5–7.5)	6–9	6–9	6–9	6–10*	6–12* (7.5–9.5*)

(continued on next page)

Table 1 (continued)

Property	<b>"<i>Natronosporangium</i>"</b>	<i>Pilimelia</i>	<i>Catellatospora</i>	<i>Catelliglobospora</i>	<i>Hamadaea</i>	<i>Actinoplanes</i>	<i>Salinispora</i>
Salt range (optimum), M of total Na <sup>+</sup>	0.1–3.0 (0.1–0.3)	nonhalophilic	nonhalophilic	nonhalophilic	nonhalophilic	Nonhalophilic; max. tolerance – 0.5 M NaCl	0–1.3 (0.1–0.3)
Temperature range (optimum)	20–40 (25)	10–38 (20–30)	15–35 (30)	20–37	4–40 (30–37)	10–35 (23–28)	10–30 (20–28)
G + C %	70.2 (genome)	70.1 (Tm)	70.7–71.4 (HPLC)	70.0 (Tm)	69.4–70.6 (HPLC)	69–73 (Tm)	69.1–70 (genome)
Habitat	Soda solonchak soil	Moderately acidic and neutral soils					Marine

\*final pH was not determined, the results were obtained on agar unbuffered media; \*\*the range is not given; PG, phosphatidylglycerol; DPG, diphosphatidylglycerol; PE, phosphatidylethanolamine; PME, phatidylmethyl ethanolamine; PI, phosphatidylinositol; PIM, phosphatidylinositolmannosyl; (V) - variable property; NR - not reported; w, weak positive.

it is highly improbable: the pH profiling was done on a solid medium, which in no way can provide stable pH conditions; the final pH was not measured; the medium was not properly buffered and, anyhow, there are no buffering systems to maintain such pH values. Furthermore, the highest reported pH maximum for bacteria measured in a pH-controlled chemostate culture is 11.3.

Since strain ACPA39<sup>T</sup> was obtained from bioprospecting polyhydrolytic haloalkaliphiles [13], its hydrolytic potential was extensively characterized by three different approaches. The cultivation-based and activity measurement-based approaches showed coherent, but slightly variable results, summarized in Table 2 and Supplementary Fig. S4. The growth experiments demonstrated that ACPA39<sup>T</sup> could utilize as growth substrate various alpha-glucans, including starch, dextrin, glycogen, cyclodextrin, pullulan and even broader range of beta-glucans, including amorphous cellulose, barley glucan, laminarin, lichenan, xylan, xyloglucan, arabinoxylan, pachyman, beta-mannan, gluco- and galactomannans, inulin. Most of them are consistent with the activity tests on plates and against dye-conjugated polysaccharides. The genome search identified a large gene inventory of glycosylhydrolases, polysaccharide lyases and carboxylesterase family enzymes (GH, PL and CE) involved in various polysaccharide hydrolysis (Table 2; Supplementary Table S2). However, in some cases, as for chitin and pectin, for example, the genome content was not supported by the negative results in growth and plate activity tests. Overall, the results demonstrated a strong polyhydrolytic potential of the novel haloalkaliphilic actinobacterium. Similar genome analysis for a presence of genes encoding the polysaccharide-specific enzymes have recently been performed in multiple available genomes of the genus *Salinispora* and yielded comparable results indicating that most of the species of these marine actinomycetes should be polyhydrolytic [47]. However, whether these actinobacteria are actually able to grow with those polysaccharides, apart from the confirmed cellulose and chitin, remains to be tested.


Among the soluble sugars, growth was observed with glucose, galactose, rhamnose, dextrose, arabinose, maltose, sucrose, trehalose, melezitose, raffinose, lactose and cellobiose. The sugars tested, but not utilized included mannose, sorbose, glucuronic and galacturonic acids, xylose, ribose and sugar alcohols mannitol and inositol.

ACPA39<sup>T</sup> showed a strong positive reaction for catalase (test with 3% H<sub>2</sub>O<sub>2</sub>) and a weak positive oxidase activity (colony assay with 0.1% tetramethyl-p-phenylenediamine HCl). Genome search identified a presence of two oxidases in ACPA39<sup>T</sup>: a *bd* quinol oxidase, often present in low oxygen-tolerant organisms, and a 4-subunit heme-copper superfamily cytochrome *c* oxidase *aa<sub>3</sub>*. But the latter was not encoded (as is usually the case) in a single genomic locus.

Ammonium and amino acids (but not nitrate or urea) can be used as the N-source. Investigation of antimicrobial activity of the isolate using disk-plate technique showed negative results with haloalkaliphilic *Bacillus horikoshii* and *Halomonas* sp. (isolated from soda lakes) and with freshwater bacteria (*E. coli*, *Pseudomonas* sp., *Bacillus subtilis*, *Micrococcus luteus* and *Staphylococcus aureus*).

#### Osmoprotection mechanism based on the genomic analysis

Since the nature of organic osmolyte was not investigated by a direct chemical analysis, an attempt was made to find the known pathways in the genome. The analysis clearly indicated that ACPA39<sup>T</sup> is using glycine betaine (GB) and related tertiary amines for osmoprotection, both importing them from environment and also by synthesizing GB *de novo*. Moreover, the genome contains a set of genes encoding a pathway of GB mineralization allowing to reutilize it as the energy and C source. The encoded import




**Fig. 2.** Maximum likelihood phylogenetic tree based on concatenated alignment of 120 conserved single copy bacterial proteins and showing position of strain ACPA39<sup>T</sup> (in bold) within the family *Micromonosporaceae*. The branch lengths correspond to the number of substitutions per site with corrections associated with the models. The black circles at nodes indicate that the percentage of corresponding support values was above 50. *Rubrobacter radiotolerans* P1<sup>T</sup> was used as an outgroup (not shown).

transporters include two ABC Opu transporters, one specific for GB (Opu AA/AB/AC) and another one with a broader tertiary amines specificity for GB/choline/carnithine import (OpuCA/CB/CD). In addition, two single-subunit GB transporters are present - OpuD and a sodium/GB symporter BetL. The imported tertiary amines can, apparently be mineralized to glycine with the released methyl groups being oxidized to CO<sub>2</sub> with the following encoded proteins identified: (1) Oxidative conversion of choline (trimethylethanol-

amine) into GB via choline dehydrogenase BetA and betaine aldehyde dehydrogenase BetB (2 copies); (2) 3-step demethylation of GB via dimethylglycine and sarcosine to glycine by a GB-homocystein methyltransferase, forming dimethylglycine and methionine, followed by dimethylglycine and sarcosine demethylation with the respective homocystein-dependent methyltransferases. Sarcosine can also be oxidized to glycine by sarcosine oxidase SoxAB. The released methyl groups are oxidized to CO<sub>2</sub>





**Fig. 3.** Influence of total Na<sup>+</sup> (as carbonates) at pH 10 (a) and pH (final values are shown) at 0.6 M total Na<sup>+</sup> (b) on growth of ACPA22<sup>T</sup> and ACPA33<sup>T</sup> on starch/yeast extract medium. The data are mean results from duplicate cultures.

via the C<sub>1</sub> pathway, including: 5,10-CH<sub>3</sub>-THF dehydrogenase Meth > methenyltetrahydrofolate dehydrogenase/cyclohydrolase F<sub>olD</sub> > 5-formyltetrahydrofolate cyclo-ligase M<sub>hs</sub> > formate dehydrogenase F<sub>doGH</sub>I. The catalytic subunits of the latter, F<sub>doGH</sub>, are located in the cytoplasm, while F<sub>doI</sub> is a di-heme b membrane subunit, which is unusual for the membrane-bound formate dehydrogenases. A potential pathway of the de novo GB biosynthesis from glycine includes three sequential methylation steps with

homocystein-dependent methyl transferases: glycine-sarcosine methyltransferase B<sub>msA</sub> (2 copies) and glycine/sarcosine/dimethylglycine N-methyltransferase, a fusion version of the two methyltransferase homologues of *Halorhodospira halophila*, which can form GB directly from glycine. A gene encoding the ectoine (a common organic osmolyte) synthase EctC is also present in the genome, but the other two subunits (EctAB) necessary for the ectoine biosynthesis are not encoded. Apart from organic osmo-

**Table 2**

Polymer-degrading potential of strain ACPA39<sup>T</sup> in growth experiments, activity tests with colony plate assay and incubations with dye-conjugated polymers and the genome-based prediction GH, glycosyl hydrolases; CE – carboxylesterases; PL – polysaccharide lyases; nd, not determined.

Polymer	Growth	Activity	CAZymes genes found in genome	
			CAZymes family	Possible function
<b>α-linked glucans</b>				
trehalose	+	+	GH15 2xGH65	trehalase trehalose phosphorylase
amylose, dextrin, cyclodextrin, pullulan glycogen	+	+	6xGH13; GH133	glycosidases with activity against alpha-linked glucans
dextran ( <i>Leuconostoc</i> )	-	-	-	dextranase
<b>β-linked polysaccharides</b>				
carboxymethyl cellulose	-	+	2x GH5; GH6; GH9	endo-beta-1,4-glucanase
amorphous cellulose	+	+	2xGH6; GH48 GH94	exoglucanase cellobiose phosphorylase
barley beta-glucan	+	+	3xGH1; 4xGH3 3xAA10	beta-glucosidase lytic cellulose monooxygenase
laminarin	+	+	2xGH16	beta-1,3-1,4 glucanase
lichenan	+	nd	3xGH1; 4xGH3	beta-glucosidase
pachyman	+	+		
xylan (beech/birch)	+	+	7xGH10; 2xGH11; 2xGH141 2xGH39; GH43	endo-beta-1,4-xylanase beta-xylosidase
			3xAA7	xylooligosaccharide oxidase
			3xCE1	acetyl xylan esterase
amorphous chitin	-	nd	2xGH18, GH23 GH3	endochitinase beta-hexosaminidase
			AA10	lytic chitin monooxygenase
			CE14	diacetylchitobiose deacetylase
<b>Other polysaccharides including heteropolysaccharides</b>				
pectin (apple, citrus)	-	-	3xPL1	pectate lyase
rhamnogalacturonan	nd	+	PL26	rhamnogalacturonan exolyase
alginate	-	nd	-	
beta-mannan	+	+	GH5; GH26	endo-beta-1,4-mannosidase
galactomannan	+	+	2x GH5; GH6; GH9	endo-beta-1,4-glucanase
glucomannan	+	nd	GH130	beta-1,4-mannooligosaccharide phosphorylase
			GH2	beta-mannosidase
			GH27; 2xGH36	alpha-galactosidase
galactan	-	-	GH2; 2xGH42	beta-galactosidase
xyloglucan	+	nd	2x GH5; GH6; GH9	endo-beta-1,4-glucanase
			3xGH1; 4xGH3	beta-glucosidase
arabinan	-	-	GH43; GH51; GH62	alpha-L-arabinosidase
arabinoxylan	+	+	7xGH10; 2xGH11; 2xGH141 GH43; GH51; GH62	endo-beta-1,4-xylanase alpha-L-arabinosidase
arabinogalactan	-	nd	-	
inulin	+	nd	GH32	inulinase

lytes, there is also a set of genes coding for potassium uptake transporters usually active in salt-tolerant microorganisms, including potassium uptake proteins TrkHA, and potassium/proton antiporters KefB and NhaP.

Overall, on the basis of the results of phylogenomic analysis and distinct phenotypic features, we propose to classify the actinobacterium isolate ACPA39<sup>T</sup> from a soda solonchak soil into a new genus and species *Natronosporangium hydrolyticum* gen. nov. sp. nov. Its comparative properties with the closely related genera of the *Micromonosporaceae* family is presented in Table 1. The common features shared with the related genera include absence of aerial mycelium; meso-DAP and lysine in the peptidoglycan; xylose as the whole cell sugar; diphosphatidylglycerol, phosphatidylethanolamine and phosphatidylinositol as the dominant membrane phospholipids; *iso*-C16:0 among the dominant PLFA and utilization of starch as growth substrate. The major difference is that ACPA39<sup>T</sup> is the only obligate alkaliphilic member of this family described so far originating from a permanently haloalkaline habitat and only a second genus (in addition to the marine genus *Salinispora*) with a moderate to high salt tolerance. Further-

more, lack of the MK-9 respiratory lipoquinones, a major domination of the *anteiso*-17:0 in the PLFA profile and the polar lipids phatidylglycerol and diphosphatidylglycerol differentiate ACPA39<sup>T</sup> from mostly related genus *Pilimelia*. Finally, ACPA39<sup>T</sup> has a remarkably broad range of polysaccharide-hydrolyzing enzymes encoded in the genome whose activity (at least partially) is confirmed in growth and activity experiments. Although this is not uncommon for actinobacteria in general, this organism can clearly be characterized as a polyhydrolytic.

The overall phylogenomic and phenotypic characteristics obtained for strain ACPA39 suggest that it can be classified in a new genus and species within the current margins of actinobacterial family *Micromonosporaceae* for which we suggest the name *Natronosporangium hydrolyticum* ge. nov., sp. nov. The results of phylogenomic analysis also indicated the necessity for a more deep phylogenetic reevaluation of the *Micromonosporaceae* family structure using the advances of the whole genome sequencing.

The new genus and species protologues are presented in Table 3.

**Table 3**

*Natronosporangium hydrolyticum* gen. nov., sp. nov.: protologue.

Parameter	Genus: <i>Natronosporangium</i> gen. nov.	Species: <i>Natronosporangium hydrolyticum</i> sp. nov.
Date created	2021-09-09	2021-09-09
Species name		<i>hydrolyticum</i>
Genus name	<i>Natronosporangium</i>	
Taxon status	gen. nov.	sp. nov.
Etymology	Na.tro.no.spo.ran'gi.um. Gr. n. <i>natron</i> , arbitrarily derived from the Arabic n. <i>natrun</i> or <i>natron</i> , soda; Gr. fam. n. <i>spora</i> a seed, a spore; Gr. neut. n. <i>angeion</i> (Latin transliteration <i>angium</i> ), vessel; N.L. neut. n. <i>Natronosporangium</i> a soda-loving organism with finger-shaped, spore-containing vessels (sporangia)	hy.dro.ly'ti.cum. Gr.neut. n. <i>hydor</i> , water; Gr. masc. adj. <i>lytikos</i> , dissolving, splitting; N.L. neut. adj. <i>hydrolyticum</i> , splitting with [by] water, referring to the hydrolytic activity of the bacterium
Description of the new taxon	Aerobic, saccharolytic, Gram-positive and filamentous actinomycetes forming mostly a well developed branched substrate mycelium with sporangia containing 2–4 spores. Peptidoglycan contains meso-DAP as the diamino acid. The membrane phospholipids include The only respiratory quinon is MK-10(H <sub>4</sub> ) and the major PLFA is a17:0. Main phenotypic features include high pH-salt tolerance and broad spectrum of hydrolyzed polysaccharides. A member of the family <i>Micromonosporaceae</i> . The type species is <i>Natronosporangium hydrolyticum</i> . The closest phylogenetic relatives are the genera <i>Pilimelia</i> and <i>Catellatospora</i> .	The aerial micelium is not developed. The substrate mycelium is highly branched, brownish in old cultures due to a formation of 2–4 spores encased in common sporangia, arranged either single or in rosettes aggregates. The spores are 0.2–0.6 μm, angular and slightly rough. Soluble pigments are not produced. The cells wall aminoacids include meso-DAP, glutamate, glycine and alanine in approximately equal proportion. The cell wall sugars are represented by arabinose, galactose and xylose. The identified membrane phospholipids include diphosphatidylglycerol, phosphatidylethanolamine, phosphatidylinositol and phosphatidylglycerol. The unidentified lipids include 4 phospho-, 4 glyco- and 2 aminophospho -lipids. The dominant PLFA include (in order of abundance) <i>anteiso</i> -C <sub>17:0</sub> , <i>iso</i> -C <sub>16:0</sub> , <i>iso</i> -C <sub>17:0</sub> , 10-Me-C <sub>18:0</sub> and <i>anteiso</i> -C <sub>17:1</sub> Ω8. The respiratory menaquinons include MK-10(H <sub>4</sub> ) (80%) and MK-11(H <sub>4</sub> ) (20%). The following sugars are utilized as substrate : glucose, galactose, rhamnose, dextrose, arabinose, maltose, sucrose, trehalose, melezitose, raffinose and lactose. The following polysaccharides are hydrolyzed: amylose, dextrin, pullulan, glycogen, pachyman, laminarin, lichenan, amorphous cellulose, carboxymethyl - and carboxyethyl-cellulose, Barley beta-glucan, xylane, pachyman, beta-mannan, glycomannan, xyloglycan, arabinoxyylan. Also hydrolyze casein and gelatin, but not olive oil. Ammonium is utilized as N-source. Oxidase is weak positive, catalase is positive. Antibiotic activity is absent. Salt-tolerant, with a range of total Na <sup>+</sup> for growth from 0.1 to 3.0 M (optimum at 0.1–0.3 M) and obligately alkaliphilic, with a pH range for growth from 7.5 to 10.5 (optimum at 8.5). At pH 10, the growth temperature range is 20–40 °C (optimum at 25). The G + C content of the DNA is 70.2% (genome). The type strain (ACPA39 <sup>T</sup> = DSM 106523 <sup>T</sup> = VKM 2772 <sup>T</sup> ) was isolated from a soda solonchak soil in the southwestern Siberia (Russia). The 16S rRNA gene sequence and the genome sequence accession numbers in the GenBank are KY775645 and CP070499.1, respectively.

(continued on next page)

Table 3 (continued)

Parameter	Genus: <i>Natronosporangium</i> gen. nov.	Species: <i>Natronosporangium hydrolyticum</i> sp. nov.
Authors (AUT)	Dimitry Y. Sorokin, Alexander I. Elcheninov, Tatiana V. Khijniak, Alicia P. Zaharycheva, Richard L. Hahnke, Olga V. Boueva, Elena V. Ariskina, Boyke Bunk, Lyudmila I. Evtushenko, Ilya V. Kublanov	
Title (TITL)	<i>Natronosporangium hydrolyticum</i> gen. nov., sp. nov., haloalkaliphilic actinobacterium from a soda solonchak soil in Central Asia	
Journal (JOUR)	Systematic and Applied Microbiology	
Corresponding author (COAU)	Dimitry Y. Sorokin	
E-mail of corresponding author (EMAU)	d.sorokin@tudelft; soroc@inmi.ru	
Designation of the type strain (TYPE)		<b>ACPA39</b>
Strain collection numbers (COLN)		DSM 106523; VKM 2772
16S rRNA gene accession number (16 SR)		KY775646
Alternative house-keeping genes: gene [accession numbers] (HKGN)		120 bacterial single-copy conserved protein markers
Genome status (GSTA)		Complete, circular chromosome (accession CP070499.1)
GC % (GGCM)		70.2 (genome)
Country of origin (COUN)	Russian Federation	Russian Federation
Region of origin (REGI)		Novosibirsk region
Date of isolation (DATI)		December 2004
Source of isolation (SOUR)	Saline alkaline soils	Soda solonchak soil
Sampling dates (DATS)		August 2003
Geographic location (GEOL)	Southwestern Siberia	Southwestern Siberia, Barabinskaya Steppe
Latitude (LATI)		54.61 N
Longitude (LONG)		76.89 E
Depth (DEPT)		0–0.1 m
Temperature of the sample (TEMS)		22 °C
pH of the sample (PHSA)		10.7
Salinity of the sample (SALS)		62 g/kg
Number of strains in study (NSTR)		1
Source of isolation of non-type strains (SAMP)		–
Growth medium, incubation conditions (CULT)	Alkaline media based on sodium carbonates, pH from 9 to 10, salinity 0.3–3 M Na <sup>+</sup>	Alkaline media based on sodium carbonates containing 0.3–0.6 M total Na <sup>+</sup> with pH 9.5–10; incubation – 37 °C; starch as C, energy and yeast extract as N-source
Conditions of preservation (PRES)	Deep freezing in 15% glycerol (v/v)	Deep freezing in 15% glycerol (v/v)
Gram stain (GRAM)	Positive	Positive
Cell shape (CSHA)	Mycelial, exospores	Moderately branched mycelium; on solid media - substrate mycelium, darkening with ageing; forms sporangia with up to 3 linearly aligned nonmotile angular exospores with smooth surface
Cell size (CSZI)		0.4–6 µm in diameter
Motility (MOTY)		nonmotile
Motility type (MOTK)		–
Type of flagellation (TFLA)		–
Sporulation (SPOR)		Exospores in sporangia
Colony morphology (COLM)		Light to dark grey, up to 2 mm; substrate mycelium only
Temperature range for growth (TEMR)		20–40 °C
Lowest temperature for growth (TEML)		20 °C
Highest temperature for growth (TEMH)	mesophilic	40
Optimal temperature for growth (TEMO)	>20	25
Lowest pH for growth (PHLO)	>7.0	7.5
Highest pH for growth (PHHI)	>9	10.3
Optimum pH for growth (PHOP)		8.5
pH category (PHCA)	Alkaliphilic	Obligate alkaliphile
Lowest salt concentration for growth (SALL)		0.1 M total Na <sup>+</sup>
Highest salt concentration for growth (SALH)		3.0 total Na <sup>+</sup>
Optimum salt concentration for growth (SALO)		0.1–0.3 total Na <sup>+</sup>
Other salts important for growth	Sodium carbonates	Sodium carbonates
Salinity category (SALC)	Halophilic	High salt-tolerant
Relation to oxygene (OREL)	Aerobic	Obligate aerobe
O <sub>2</sub> conditions for strain testing (OCON)		Air
Carbon source used (class) (CSUC)	Carbohydrates	Carbohydrates, proteins
Specific compounds (CSUC)	Polysaccharides (alpha/beta-glucans)	<b>Sugars:</b> cellobiose. <b>Polysaccharides:</b> cyclodextrin, glycogen, pullulan, inulin, amorphous cellulose, laminarin, lichenan, barley glucan, xylane, pachyman, beta-mannan, glucomannan, xyloglycan, arabinoxylan.
Nitrogen source (NSOU)		Ammonium, yeast extract
Terminal electron acceptor (ELAC)	O <sub>2</sub>	O <sub>2</sub>
Energy metabolism (EMET)	chemoorganoheterophilic	chemoorganoheterophilic
Phospholipids (PHOS)		PG, DPG, PE, PI, PIM
Glycolipids (GLYC)	Glyco- and aminoglycolipids	5 unidentified glycolipids
Respiratory quinons	MK-10	MK-10(H4)
Habitat (HABT)	Saline alkaline soils	Soda solonchak soil, s-w Siberia
Extraordinary feautres (EXTR)	Polyhydrolytics	Utilize large number of alpha/beta-glucans for growth at haloalkaline conditions

The 16S rRNA gene and whole genome sequences of strains ACPA39<sup>T</sup> have been deposited in the NCBI GenBank under the numbers KY775646 and CP070499.1, respectively. Raw read data were deposited at NCBI SRA (698925).

## Acknowledgements

The authors thank Nicole Heyer and Bettina Ehlert for excellent technical assistance.

## Funding information

This work was supported in part (isolation and phenotypical characterization) by the Russian Science Foundation (20-14-00250) to DYS, AGE and IVK, and by the Russian Ministry of Science and Higher Education (genome annotation and analysis, phylogeny, electron microscopy) to DYS, AGE, IVK and LIE.

## Appendix A. Supplementary data

Supplementary data to this article can be found online at <https://doi.org/10.1016/j.syapm.2022.126307>.

## References

- [1] Kroppenstedt, R.M., Evtushenko, L.I. (2006) The family *Nocardiopsaceae*. *Prokaryotes* 3, 754–795.
- [2] Shivilata, L., Satyanarayana, T. (2015) Thermophilic and alkaliphilic *Actinobacteria*: biology and potential applications. *Front. Microbiol.* 6, 1014.
- [3] Antony-Babu, S., Goodfellow, M. (2008) Biosystematics of alkaliphilic streptomycetes isolated from seven locations across a beach and dune sand system. *Ant van Leeuwenhoek* 94 (4), 581–591.
- [4] Grant, W.D., Jones, B.E. Bacteria, archaea and viruses of soda lakes. In: Schagerl, M. (Ed.), *Soda Lakes of East Africa*. Springer International Publishing, Cham, pp. 97–147. [https://doi.org/10.1007/978-3-319-28622-8\\_5](https://doi.org/10.1007/978-3-319-28622-8_5).
- [5] Kondorskaya, N.I. (1965) Geographic distribution of soda soils in USSR. *Soil Sci. (Moscow, in Russian)* No 9, 10–16.
- [6] Bazilevich, N.I. (1970) The geochemistry of soda soils. USDA, NSF and Israel Program for Scientific Translations. Jerusalem, 396 p.
- [7] Yang, R., Zhang, L.-P., Guo, L.-G., Shi, N., Lu, Z., Zhang, X. (2008) *Nocardiopsis valliformis* sp. nov., an alkaliphilic actinomycete isolated from alkali lake soil in China. *Int. J. Syst. Evol. Microbiol.* 58 (7), 1542–1546.
- [8] Mwirichia, R., Muigai, A.W., Tindall, B., Boga, H.I., Stackebrandt, E. (2010) Isolation and characterisation of bacteria from the haloalkaline Lake Elmenteita, Kenya. *Extremophiles* 14 (4), 339–348.
- [9] Ara, I., Daram, D., Baljinova, T., Yamamura, H., Hozzein, W.N., Bakir, M.A. (2013) Isolation, classification, phylogenetic analysis and scanning electron microscopy of halophilic, halotolerant and alkaliphilic actinomycetes isolated from hypersaline soil. *Afr. J. Microbiol. Res.* 7, 298–308.
- [10] Akhwale, J.K., Göker, M., Rohde, M., Spröer, C., Schumann, P., Klenk, H.-P., Boga, H.I. (2015) *Streptomyces alkaliphilus* sp. nov., isolated from sediments of Lake Elmenteita in the Kenyan Rift Valley. *Ant van Leeuwenhoek* 107 (5), 1249–1259.
- [11] Sorokin, D.Y. (2005) Is there a limit for high-pH growth? *Int. J. Syst. Evol. Microbiol.* 55, 1405–1406.
- [12] Maldonado, L.A., Fenical, W., Jensen, P.R., Kauffman, C.A., Mincer, T.J., Ward, A.C., Bull, A.T., Goodfellow, M. (2005) *Salinispora arenicola* gen. nov., sp. nov. and *Salinispora tropica* sp. nov., obligate marine actinomycetes belonging to the family *Micromonosporaceae*. *Int. J. Syst. Evol. Microbiol.* 5, 1759–1766.
- [13] Sorokin, D.Y., Khijniak, T.V., Kolganova, T.V., Jones, B.E., Kublanov, I.V. (2017) Culturable diversity of aerobic polyhydrolytic haloalkaliphilic bacteria in saline alkaline soils. *Peer J.* 5, e3796.
- [14] Sorokin, D.Y., Khijniak, T.V., Zaharycheva, A.P., Elcheninov, A.G., Hahnke, R.L., Boueva, O.V., Ariskina, E.V., Bunk, B., Kublanov, I.V., Evtushenko, L.I. (2021) *Natronoglycomyces albus* gen. nov., sp. nov., a haloalkaliphilic actinobacterium from a soda solonchak soil. *Int. J. Syst. Evol. Microbiol.* 71, 004804.
- [15] Shirling, E.B., Gottlieb, D. (1966) Methods for characterization of *Streptomyces* species. *Int. J. Syst. Bacteriol.* 16 (3), 313–340.
- [16] Reasoner, D.J., Geldreich, E.E. (1985) A new medium for the enumeration and subculture of bacteria from potable water. *Appl. Environ. Microbiol.* 49 (1), 1–7.
- [17] Boone, C.J., Pine, L. (1968) Rapid method for characterization of actinomycetes by cell wall composition. *Appl. Microbiol.* 16 (2), 279–284.
- [18] Moore, S., Stein, W.H. (1969) Chromatographic determination of amino acids by the use of automatic recording equipment. *Methods Enzymol.* 6, 819–831.
- [19] Sinner, M., Puls, J. (1978) Non-corrosive dye reagent for detection of reducing sugars in borate complex ion-exchange chromatography. *J. Chromatogr. A* 156 (1), 197–204.
- [20] Minnikin, D.E., O'Donnell, A.G., Goodfellow, M., Alderson, G., Athalye, M., Schaal, A., Parlett, J.H. (1984) An integrated procedure for the extraction of bacterial isoprenoid quinones and polar lipids. *J. Microbiol. Methods* 2 (5), 233–241.
- [21] Kudryashova, E.B., Karlyshev, A.V., Ariskina, E.V., Streshinskaya, G.M., Vinokurova, N.G., Koptsyn, D.S., Evtushenko, L.I. (2018) *Cohnella kolyemensis* sp. nov., a novel bacillus isolated from Siberian permafrost. *Int. J. Syst. Evol. Microbiol.* 68, 2912–2917.
- [22] Collins, M.D. (1985) Analysis of isoprenoid quinones. *Meth. Microbiol.* 18, 329–363.
- [23] Zhang, H., Yohe, T., Huang, L.e., Entwistle, S., Wu, P., Yang, Z., Busk, P.K., Xu, Y., Yin, Y. (2018) dbCAN2: a meta server for automated carbohydrate-active enzyme annotation. *Nucleic Acids Res.* 46 (W1), W95–W101.
- [24] Park, D. (2007) Genomic DNA isolation from different biological materials. *Meth. Mol. Biol.* 353, 3–13.
- [25] Katoh, K., Misawa, K., Kum, K., Miyata, T. (2002) MAFFT: a novel method for rapid multiple sequence alignment based on fast Fourier transform. *Nucleic Acids Res.* 30, 3059–3066.
- [26] Kumar, S., Stecher, G., Tamura, K. (2016) MEGA7: Molecular Evolutionary Genetics Analysis Version 7.0 for Bigger Datasets. *Mol. Biol. Evol.* 33 (7), 1870–1874. <https://doi.org/10.1093/molbev/msw054>.
- [27] Parks, D.H., Rinke, C., Chuvochina, M., Chaumeil, P.-A., Woodcroft, B.J., Evans, P.N., Hugenholtz, P., Tyson, G.W. (2017) Recovery of nearly 8,000 metagenome-assembled genomes substantially expands the tree of life. *Nat. Microbiol.* 2 (11), 1533–1542.
- [28] Parks, D.H., Chuvochina, M., Waite, D.W., Rinke, C., Skarshewski, A., Chaumeil, P.-A., Hugenholtz, P. (2018) A standardized bacterial taxonomy based on genome phylogeny substantially revises the tree of life. *Nat. Biotechnol.* 36 (10), 996–1004.
- [29] Capella-Gutierrez, S., Silla-Martinez, J.M., Gabaldon, T. (2009) trimAl: A tool for automated alignment trimming in large-scale phylogenetic analyses. *Bioinformatics* 25 (15), 1972–1973.
- [30] Stamatakis, A. (2014) RAxML version 8: A tool for phylogenetic analysis and post-analysis of large phylogenies. *Bioinformatics* 30 (9), 1312–1313.
- [31] Letunic, I., Bork, P. (2019) Interactive Tree Of Life (iTOL) v4: recent updates and new developments. *Nucleic Acids Res.* 47, W256–W259.
- [32] Pritchard, L., Glover, R.H., Humphris, S., Elphinstone, J.G., Toth, I.K. (2016) Genomics and taxonomy in diagnostics for food security: Soft-rotting enterobacterial plant pathogens. *Anal. Methods* 8, 12–24.
- [33] Rodríguez-R, L., Konstantinidis, K. (2016) The enveomics collection: a toolbox for specialized analyses of microbial genomes and metagenomes. *Peer J. Prepr.* <https://doi.org/10.7287/peerj.preprints.1900>.
- [34] Buchfink, B., Xie, C., Huson, D.H. (2015) Fast and sensitive protein alignment using DIAMOND. *Nat. Methods* 12 (1), 59–60.
- [35] Collins, M.D., Jones, D. (1981) Distribution of isoprenoid quinone structural types in bacteria and their taxonomic implication. *Microbiol. Rev.* 45 (2), 316–354.
- [36] Genilloud O. (2015) *Micromonosporaceae*. In: Whitman, W.B. (Ed.), *Bergey's Manual of Systematic of Bacteria and Archaea*. John Wiley & Sons. doi: 10.1002/9781118960608.fbm00148.
- [37] Salam, N., Jiao, J.-Y., Zhang, X.-T., Li, W.-J. (2020) Update on the classification of higher ranks in the phylum *Actinobacteria*. *Int. J. Syst. Evol. Microbiol.* 70 (2), 1331–1355.
- [38] Konstantinidis, K.T., Rosselló-Móra, R., Amann, R. (2017) Uncultivated microbes in need of their own taxonomy. *ISME J.* 11 (11), 2399–2406.
- [39] Qin, Q.-L., Xie, B.-B., Zhang, X.-Y., Chen, X.-L., Zhou, B.-C., Zhou, J., Oren, A., Zhang, Y.-Z. (2014) A proposed genus boundary for the prokaryotes based on genomic insights. *J. Bacteriol.* 196 (12), 2210–2215.
- [40] Aliyu, H., Lebre, P., Blom, J., Cowan, D., De Maayer, P. (2016) Phylogenomic re-assessment of the thermophilic genus *Geobacillus*. *Syst. Appl. Microbiol.* 39 (8), 527–533.
- [41] Lopes-Santos, L., Castro, D.B.A., Ferreira-Tonin, M., Corrêa, D.B.A., Weir, B.S., Park, D., Ottoboni, L.M.M., Neto, J.R., Destéfano, S.A.L. (2017) Reassessment of the taxonomic position of *Burkholderia andropogonis* and description of *Robbsia andropogonis* gen. nov., comb. nov. *Antonie Van Leeuwenhoek* 110 (6), 727–736.
- [42] Li, Y., Xue, H., Sang, S.-Q., Lin, C.-L., Wang, X.-Z. (2017) Phylogenetic analysis of family Neisseriaceae based on genome sequences and description of *Populibacter corticis* gen. nov., sp. nov., a member of the family Neisseriaceae, isolated from symptomatic bark of *Populus euramericana* canker. *PLoS ONE* 12, e0174506.
- [43] Wirth, J.S., Whitman, W.B. (2018) Phylogenomic analyses of a clade within the roseobacter group suggest taxonomic reassignments of species of the genera *Aestuuarivita*, *Citricella*, *Loktanelia*, *Nautella*, *Pelagibaca*, *Ruegeria*, *Thalassobius*, *Thiobacimonas* and *Tropicibacter*, and the proposal of six novel genera. *Int. J. Syst. Evol. Microbiol.* 68, 2393–2411.
- [44] Orata, F.D., Meier-Kolthoff, J.P., Sauvageau, D., Stein, L.Y. (2018) Phylogenomic analysis of the gammaproteobacterial methanotrophs (order *Methyloccales*) calls for the reclassification of members at the genus and species levels. *Front. Microbiol.* 9, 3162.
- [45] Xu, Z., Masuda, Y., Itoh, H., Ushijima, N., Shiratori, Y., Senoo, K. (2019) *Geomonas oryzae* gen. nov., sp. nov., *Geomonas edaphica* sp. nov., *Geomonas ferrireducens* sp. nov., *Geomonas terrae* sp. nov., four ferric-reducing bacteria isolated from paddy soil, and reclassification of three species of the genus *Geobacter* as members of the genus *Geomonas* gen. nov. *Front. Microbiol.* 10, 2201.

- [46] Ito, M., Morino, M., Krulwich, T.A. (2017) Mrp antiporters have important roles in diverse bacteria and archaea. *Front. Microbiol.* 8, 2325.
- [47] Roman-Ponce, B., Millan-Aguinaga, N., Guillen-Matus, D., Chase, A.B., Ginigini, J.G.M., Soapi, K., Feussner, K.D., Jensen, P.R., Trujillo, M.E. (2020) Six novel species of the obligate marine actinobacterium *Salinispora*, *Salinispora cortesiana* sp. nov., *Salinispora fenicalii* sp. nov., *Salinispora goodfellowii* sp. nov., *Salinispora mooreana* sp. nov., *Salinispora oceanensis* sp. nov. and *Salinispora vitiensis* sp. nov., and emended description of the genus *Salinispora*. *Int. J. Syst. Evol. Microbiol.* 70, 4668–4682.
- [48] Ara, I., Bakir, M.A., Kudo, T. (2008) Transfer of *Catellatospora koreensis* Lee et al. 2000 as *Catelliglobospora koreensis* gen. nov., comb. nov. and *Catellatospora tsunoense* Asano et al. 1989 as *Hamadaea tsunoensis* gen. nov., comb. nov., and emended description of the genus *Catellatospora* Asano and Kawamoto 1986 emend. Lee and Hah 2002. *Int. J. Syst. Evol. Microbiol.* 58, 1950–1960.
- [49] Vobis, G., Kömpfer, P. (2015) *Pilimelia*. In: Whitman, W.B. (Ed.), *Bergey's Manual of Systematic of Bacteria and Archaea*. John Wiley & Sons. doi: 10.1002/9781118960608.gbm00149.
- [50] Raney, F. (2015) *Catellatospora*. In: Whitman, W.B. (Ed.), *Bergey's Manual of Systematic of Bacteria and Archaea*. John Wiley & Sons. doi: 10.1002/9781118960608.gbm00141.
- [51] Chu, X., Li, S., Chen, W., Asem, M.D., Duan, Y.-Q., Nie, G.-X., Hozzein, W.N., Zhi, X.-Y., Li, W.-J. (2016) *Hamadaea flava* sp. nov., isolated from a soil sample and emended description of the genus *Hamadaea*. *Int. J. Syst. Evol. Microbiol.* 66, 1818–1822.
- [52] Sun, Y., Jiang Z.-M., Zhao, L.-L., Su J., Yu, L.-Y., Tian, Y.-Q., Zhang, Y.-Q. *Allorhizocola rhizosphaerae* gen. nov., sp. nov., a new member of *Micromonosporaceae* isolated from rhizosphere soil of the plant *Calligonum mongolicum*. *Int. J. Syst. Evol. Microbiol.* 69, 109–115.
- [53] Vobis, G., Schäfer, J., Kömpfer, P. (2015) *Actinoplanes*. In: Whitman, W.B. (Ed.), *Bergey's Manual of Systematic of Bacteria and Archaea*. John Wiley & Sons. doi: 10.1002/9781118960608.gbm00139.