



Delft University of Technology

Document Version

Final published version

Licence

CC BY

Citation (APA)

Wang, Y., Lan, D., Durrani, R., & Hollmann, F. (2017). Peroxygenases en route to becoming dream catalysts. What are the opportunities and challenges? *Current Opinion in Chemical Biology*, 37, 1-9.
<https://doi.org/10.1016/j.cbpa.2016.10.007>

Important note

To cite this publication, please use the final published version (if applicable).
Please check the document version above.

Copyright

In case the licence states "Dutch Copyright Act (Article 25fa)", this publication was made available Green Open Access via the TU Delft Institutional Repository pursuant to Dutch Copyright Act (Article 25fa, the Taverne amendment). This provision does not affect copyright ownership.

Unless copyright is transferred by contract or statute, it remains with the copyright holder.

Sharing and reuse

Other than for strictly personal use, it is not permitted to download, forward or distribute the text or part of it, without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license such as Creative Commons.

Takedown policy

Please contact us and provide details if you believe this document breaches copyrights.
We will remove access to the work immediately and investigate your claim.

This work is downloaded from Delft University of Technology.

Peroxygenases *en route* to becoming dream catalysts. What are the opportunities and challenges?

Yonghua Wang¹, Dongming Lan¹, Rabia Durrani² and
Frank Hollmann³



Peroxygenases are promising catalysts for preparative oxyfunctionalization chemistry as they combine the versatility of P450 monooxygenases with simplicity of cofactor-independent enzymes. Though many interesting applications have been reported, today 'we have only scratched the surface' and significant efforts are necessary to solve issues related to selectivity of the wild type enzymes and low product titers. For this, further elucidation of the vast natural diversity as well as protein and reaction engineering approaches are discussed.

Addresses

¹ School of Food Science and Engineering, South China University of Technology, Guangzhou 510640, China
² School of Bioscience and Bioengineering, South China University of Technology, Guangzhou 510006, China
³ Department of Biotechnology, Delft University of Technology, Van der Maasweg 9, 2629HZ Delft, The Netherlands

Corresponding author: Hollmann, Frank (F.Hollmann@tudelft.nl)

Current Opinion in Chemical Biology 2017, 37:1–9

This review comes from a themed issue on **Biocatalysis and Biotransformation**

Edited by **Bernhard Hauer** and **Stefan Lutz**

For a complete overview see the [Issue](#) and the [Editorial](#)

Available online 16th December 2016

<http://dx.doi.org/10.1016/j.cbpa.2016.10.007>

1367-5931/© 2016 The Authors. Published by Elsevier Ltd. This is an open access article under the CC BY license (<http://creativecommons.org/licenses/by/4.0/>).

Introduction

The selective activation of inert or poorly activated C–H bonds certainly is a 'dream reaction' of organic chemistry. Today transition metals are the catalysts of choice for the insertion of activated oxygen [1]. However, the selectivity of low-molecular weight-catalysts often is dictated by the intrinsic properties of the starting material (i.e. bond dissociation energies and steric constraints) and therefore offers little possibilities to control the selectivity of the oxyfunctionalization reaction. However, when embedded into a well-defined cavity (such as in proteins), selectivity can be imposed by the supramolecular 'ligand' thereby overriding the chemical reactivity of the starting material. Especially heme-thiolate containing enzymes have been investigated thoroughly in the past years amongst them the well-known P450 monooxygenases [2–4] and, more recently, peroxygenases (E.C. 1.11.2.1) [5*,6]. Both enzyme

classes rely on oxoferryl-heme as the oxygenating species (Compound I) to catalyze a broad range of oxyfunctionalization reactions (Scheme 1).

In P450 monooxygenases Compound I is regenerated through a sequence of reductive activation of molecular oxygen involving reduced nicotinamide cofactors and more or less complicated multi-enzyme electron transport chains [7,8]. Peroxygenases utilize partially reduced oxygen (H_2O_2) directly (hydrogen peroxide shunt pathway). The border between both enzyme classes is sometimes fluent as so-called P450 peroxygenases are capable of utilizing both pathways [9].

From an organic chemistry point-of-view the simplicity of peroxygenases is appealing. Since the first report on a novel peroxygenases from *Agrocybe aegerita* in 2004 [10] the last decade has seen a considerably increasing interest in peroxygenases [5*,6]. The aim of this contribution is to critically summarize — from a chemist's point-of-view — the most relevant developments and identify current bottlenecks together with promising solutions.

Structure and mechanism

Today, only two peroxygenase crystal structures have been published [11*,12]. The overall structure of the peroxygenase from *Agrocybe aegerita* (AaeUPO) is shown in Figure 1.


The catalytic mechanism of AaeUPO has been discussed in detail by Hofrichter and coworkers [5*]. Here, it is worth pointing out the role of the distal glutamate 196 and arginine 189 involved through acid–base catalysis in the H_2O_2 -activation step (Figure 1); the homologous *Cfu*UPO differs in this respect (histidine instead of arginine), which may also account for the sometimes dramatic differences in reactivity between both enzymes.

Overall, peroxygenases allow, in principle, for the same, rich oxyfunctionalization chemistry as the P450 monooxygenases while being independent from reduced nicotinamide cofactors and complicated electron transport chains. Particularly this feature makes peroxygenases interesting catalysts for organic synthesis.

Applications of peroxygenases

Several dozen different reactions including hydroxylation of (non-)activated C–H bonds, epoxidations, heteroatom

Scheme 1



Compound I as the active catalyst within P450 monooxygenases and peroxygenases to perform selective oxyfunctionalization reactions. While P450 monooxygenases regenerate Cpd I via sequential reduction of molecular oxygen peroxygenases (shown here) form Cpd I directly from hydrogen peroxide.

oxygenations etc. have been reported. A more extensive discussion of the details can be found in two recent review articles [5[•],6]. Here, we would like to critically evaluate the current state-of-the-art focusing on selectivity issues in particular.

Alkane hydroxylation


Aliphatic substrates ranging from ethane to fatty acids can be hydroxylated by peroxygenases [5[•],6]. As shown in Figure 2 the product distribution and selectivity observed in these reactions can significantly vary with the biocatalyst used but also by the substrate properties. For example, fatty acids are converted to a mixture of ω , ω -1 and ω -2 hydroxylation products using *AaeUPO* (Figure 2b) [13]. Likewise, linear alkanes preferentially yield a mixture of 2-alkanols and 3-alkanols [14]. Quite remarkably, while the regioselectivity is comparably poor, the enantioselectivity can be high. A completely different regioselectivity

is observed when converting fatty acids with the P450 peroxygenases from *Bacillus subtilis* (P450_{Bs β}) or *Clostridium acetobutylicum* (P450_{Cl α}) [15,16]. With these enzymes α -hydroxylation or β -hydroxylation is observed, respectively (Figure 2c).

In a recent contribution, Gutiérrez and coworkers showed that the hydroxylation of cholecalciferol (Vitamin D) when catalyzed by *AaeUPO* is rather unselective whereas the peroxygenase from *Coprinopsis cinerea* (*CciUPO*) essentially gave only one product (Figure 2d). The authors rationalized this observation by differences in the size of the substrate access channels and different degrees of translational freedom of the substrates [17[•]].


Another issue of peroxygenase-catalyzed hydroxylation of alkanes is the frequently observed subsequent conversion of the alcohols into the corresponding aldehydes and

Figure 1



Overall structure of AaeUPO (2YOR, upper) and its proposed catalytic mechanism (lower). In the first step H_2O_2 displaces water as the 6th Fe ligand. The resulting peroxocomplex is deprotonated by Glu196, which also facilitates the heterolytic cleavage of the O-O-bond resulting in the active Compound I (Cpd I). The latter performs a H-atom abstraction at the substrate (R-H) leaving an enzyme-bound radical which swiftly recombines with the hydroxy ligand. After dissociation of the product a new catalytic cycle begins.

Figure 2



Current Opinion in Chemical Biology

Selected examples for alkane hydroxylation/alkene epoxidation focusing on selectivity issues. **(a)** Alkane hydroxylation (e.g. of cyclohexane) is frequently plagued by undesired overoxidation to the corresponding aldehydes and ketones [18]; **(b)** alkane hydroxylation (e.g. within fatty acids) generally leads to a mixture of predominantly ω -1-hydroxylation and ω -2-hydroxylation products [13]; **(c)** in contrast, some P450 peroxygenases also catalyze α -hydroxylation and β -hydroxylation of fatty acids [15,16]; **(d)** the selectivity of alkane hydroxylation (e.g. with Vitamin D) can significantly vary between different peroxygenases [17*]; **(e)** epoxidation generally competes with allylic hydroxylation [19]; **(f)** aromatic hydroxylation (especially in the absence of directing groups) can lead to different regioisomers and — more importantly — also further oxidation and radical polymerization of the resulting phenol products [20].

ketones (Figure 2a) [18]. Especially if the (chiral) alcohol is the product of interest, this ‘overoxidation’ is highly undesirable. Again, the extent of this ‘overoxidation’ reaction can vary between peroxygenases from different sources: the peroxygenase from *Marasmius rotula* (*Mro*UPO) catalyzes the oxidation of cyclohexanol to cyclohexanone quite efficiently performed while *Aae*UPO and *Cci*UPO show significantly reduced overoxidation activities [18]. As a result, selective accumulation of either cyclohexanol or cyclohexanone starting from cyclohexane could be achieved.

Substrates with activated C–H bonds are often converted more selectively. For example, ethers and secondary amines are generally attacked at α -position to the heteroatom leading to dealkylation reactions [21].

*Aae*UPO-catalyzed hydroxylations of benzylic C–H bonds occur highly regioselectively and enantioselectively [22]. However, with an increasing steric demand of the alkyl sidechain a decreasing enantioselectivity was observed. Again, overoxidation generally represents an undesired side-reaction.

Aromatic hydroxylation

Various aromatic hydroxylations have been reported using peroxygenases. Amongst them the regioselective hydroxylation of 2-(4-hydroxyphenoxy)propionic acid [23] and hydroxypropranolol or diclofenac [24]. The mechanism of arene hydroxylation involves epoxide intermediates, which spontaneously rearrange to the corresponding phenol (Figure 2f) [20,25]. Typically, the intermediate epoxides are released into the reaction mixture suggesting that the biocatalysts is not directly involved in the rearrangement reaction leading to the phenol product.

A challenge with arene hydroxylation is that the phenol products often undergo peroxygenase-catalyzed H-atom abstraction yielding phenoxy radicals, which spontaneously polymerize (Figure 2f) [26,27]. Generally, this is avoided by application of radical scavengers such as ascorbic acid [26]. Alternatively, protein engineering has been shown to efficiently circumvent this [20].

Epoxidations

As shown in Figure 2e, peroxygenase-catalyzed epoxidation reactions are generally plagued by a comparably poor chemoselectivity. Because of their relative lability, allylic C–H-bonds are most frequently hydroxylated as well yielding complex product mixtures [22,28,29]. Allylic methyl groups appear to be less reactive as compared to methylene groups. Hence, selective epoxidation is observed with sub-terminal alkenes such as methyl styrenes. Particularly α -substituted and β -substituted styrenes appear to be converted with high enantioselectivity whereas other styrene derivates yield near-racemic epoxides.

Miscellaneous reactions

Next to the ‘classical’ oxyfunctionalization reactions mentioned above, the recent years have also seen some new applications worth to be shortly discussed here.

Recently, oxidative decarboxylation of carboxylic acids yielding terminal olefins has been reported with OleT (a P450 monooxygenase from *Jeotgalicoccus* sp. ATCC 8456 exhibiting significant peroxygenase activity), which may become interesting in view of transforming renewable materials into chemical building blocks (Figure 3a) [30*,31–33].

A very interesting novel application of *Cfu*UPO has been reported by Deska and coworkers (Figure 3b) [34**]. The authors demonstrated that *Cfu*UPO also catalyzes the conversion of furylcarbinols (obtained from enantioselective ADH-catalyzed reduction of the prochiral ketone precursors) to pyranones (Achmatowicz reaction). Because of the importance of this reaction in natural product synthesis further exciting developments may be expected here.

Also the, long-neglected haloperoxidase activity of many peroxygenases is receiving a renewed interest. Here, the ability of peroxidases/peroxygenases to generate hypohalous acids, which undergo spontaneous, non-enzymatic electrophilic oxidation reactions is exploited [36,37*,39–43]. From a green chemistry point of view, the avoidance of stoichiometric amounts of bleach together with the resulting salt wastes is of interest. For example, Holtmann and coworkers reported an electroenzymatic system for the chlorination of thymol (Figure 3c) [36].


The above-mentioned overoxidation is not always undesired as for example in case of the conversion of hydroxymethyl furfural to furanedicarboxylic acid (Figure 3d) [38]. The combination of a H_2O_2 -generating oxidase with (H_2O_2 -dependent) *Aae*UPO enabled a ‘self-sufficient’, aerobic oxidation reaction.

Overall it can be concluded that the selectivity of peroxygenase-catalyzed reactions often is dominated by the intrinsic reactivity of the substrates and highly selective reactions occur ‘by chance’. Turning peroxygenases into highly selective catalysts will be one of the major tasks for future development as only highly selective peroxygenases will be truly practical catalyst for organic chemistry.

Improved peroxygenases from natural or man-made diversity

Today, more than 1000 putative peroxygenase genes are deposited in genome databases [5*]. Only a fraction of these peroxygenases have been elucidated yet with respect to their as potential catalysts for chemical synthesis. We are certain that the near future will bring about exciting new enzymes!

Figure 3



Selection of new, upcoming oxidative transformations utilizing peroxygenases. **(a)** Oxidative decarboxylation of (fatty) acids yielding terminal olefins [30[•],31[•]–33], **(b)** biocatalytic Achmatowicz reactions [34^{••},35], **(c)** electrophilic halogenation reactions [36,37[•]], **(d)** oxidase/peroxygenase cascades for example for the transformation of HMF into furan dicarboxylic acid [38].

However, also *AaeUPO* is an excellent starting point for improved peroxygenases. However, to fully exploit its catalytic potential, access to mutants with tailored properties is mandatory [44,45]. To generate and select improved enzymes the following tools are necessary: (1) an efficient expression system, (2) a reliable screening assay and (3) a smart method to generate a mutants libraries.

As a glycoprotein, functional expression of the *AaeUPO* is not straightforward in *Escherichia coli*, which is why current research efforts focus on fungal expression systems. Alcalde and coworkers succeeded in the expression of *AaeUPO* in *Saccharomyces cerevisiae* [46^{••},47[•]]. Particularly, evolution of the signal peptide led to a dramatic improvement the enzyme titer from originally 0.007 mg/L to 217 mg/L in *Pichia pastoris*.

In order to screen large mutants libraries, fast and reliable screening assays are required. Especially, photometric assays are suitable due to the sensitivity and applicability in high throughput format. For peroxygenases different

assays detecting peroxidase activity, peroxygenase activity and haloperoxidase activity have been reported [9]. Alcalde and coworkers for example used a smart combination of two different assays to simultaneously increase *AaeUPO*'s arene hydroxylation activity and decrease its phenol polymerization activity [20].

The aforementioned assay is suitable for general enzyme properties such as activity and stability. However, if modification of the product scope (selectivity) is desired, more specific assays are necessary. For this chromatographic assays are state of the art but are more time consuming than simple spectrophotometric assays. Therefore, reducing the actual library size is of utmost importance. Fortunately, the crystal structure of *AaeUPO* is available [11[•]]. This will enable the generation of smart and high quality focused libraries. Exciting developments can be expected in the near future.

Reaction engineering

Two further bottlenecks *en route* to preparative application of peroxygenases are worth mentioning here. First,

peroxygenases, just like all heme-dependent enzymes, are rather sensitive against H_2O_2 [48]. Therefore, maintaining the H_2O_2 concentration at an optimal level high enough to sustain the peroxygenases reaction and low enough to avoid oxidative inactivation is mandatory. The most promising approach is *in situ* H_2O_2 generation through reductive activation of O_2 . Various approaches have been proposed to control the H_2O_2 formation rate by adjusting the catalyst concentration. Today, the system glucose oxidase prevails due to its simplicity and the cheap, commercially available reagents [49]. One major disadvantage (especially envisioning large-scale applications) however is its poor atom-efficiency and the stoichiometric accumulation of gluconolactone (or the corresponding gluconic acid). Electrochemical methods [36,50,51] implying cathodic reduction of molecular oxygen are principally better suited but necessitate electrochemical equipment. The same is true for the proposed photochemical approaches [52,53]. Recently, we proposed an enzyme cascade enabling the complete oxidation of methanol to CO_2 and productive use of all reducing equivalents liberated in the oxidation steps to promote peroxygenases-catalyzed hydroxylations [54^{**}].

Secondly, the rather low substrate loadings reported with peroxygenases so far has to be improved significantly! Typically, 1–10 mM of starting material is converted, corresponding to a maximal product titer of less than 1 g L^{-1} which obviously impairs the synthetic attractiveness of the current reaction schemes [55[•],56[•]]. A range of interesting reaction engineering approaches have been proposed to overcome this limitation, amongst them the application of the so-called two-liquid-phase approach. Here, a hydrophobic organic phase (ideally the substrate itself) serves as a substrate reservoir and as product sink [57]. Neat reaction conditions, that is the avoidance of additional solvents would be most efficient [58].

Future developments

Peroxygenases have the potential of becoming very useful tools for organic oxyfunctionalization chemistry. Today's catalysts suffer from poor selectivity necessitating a significantly enlarged diversity of enzymes to choose from. On the one hand, the large natural diversity of peroxygenases still remains to be explored. We believe that today we have just scratched the surface of what nature has to offer to the organic chemist. On the other hand, man-made diversity obtained from random or (semi-)rational protein engineering has proven to deliver tailor-made enzymes [45,59–61]. We are convinced that applying the knowledge and techniques developed in this area during the past 20 years [62] will provide the chemist with selective and robust peroxygenases.

Another challenge of UPOs will be the poor water solubility of most reagents of interest. Today, this is met using

water miscible cosolvents such as acetonitrile or acetone. We, however, believe that these solutions will not be practical especially for large-scale applications as the substrate loadings here mostly remain below a few grams per liter. Alternative reaction concepts such as the two-liquid-phase approach or even neat reaction systems will be necessary to evoke the interest of potential users.

Acknowledgements

The work is supported by National Natural Science Foundation of China (31471690) and by the European Research Council (ERC) through the ERC Consolidator Grant (No 648026).

References and recommended reading

Papers of particular interest, published within the period of review, have been highlighted as:

- of special interest
- of outstanding interest

1. Roduner E, Kaim W, Sarkar B, Urlacher VB, Pleiss J, Gläser R, Einicke W-D, Sprenger GA, Beifuß U, Klemm E et al.: **Selective catalytic oxidation of C–H bonds with molecular oxygen.** *ChemCatChem* 2013, **5**:82–112.
2. Fasan R: **Tuning P450 enzymes as oxidation catalysts.** *ACS Catal* 2012, **2**:647–666.
3. Urlacher VB, Girhard M: **Cytochrome P450 monooxygenases: an update on perspectives for synthetic application.** *Trends Biotechnol* 2012, **30**:26–36.
4. Bernhardt R, Urlacher VB: **Cytochromes P450 as promising catalysts for biotechnological application: chances and limitations.** *Appl Microbiol Biotechnol* 2014, **98**:6185–6203.
5. Hofrichter M, Ullrich R: **Oxidations catalyzed by fungal peroxygenases.** *Curr Opin Chem Biol* 2014, **19**:116–125.

This recent review nicely summarizes the chemical transformations accessible through peroxygenases.

6. Bormann S, Gomez Baraibar A, Ni Y, Holtmann D, Hollmann F: **Specific oxyfunctionalisations catalysed by peroxygenases: opportunities, challenges and solutions.** *Catal Sci Technol* 2015, **5**:2038–2052.
7. Holtmann D, Fraaije MW, Opperman DJ, Arends IWCE, Hollmann F: **The taming of oxygen: biocatalytic oxyfunctionalisations.** *Chem Commun* 2014, **50**:13180–13200.
8. Holtmann D, Hollmann F: **The oxygen dilemma: a severe challenge for the application of monooxygenases?** *ChemBioChem* 2016, **17**:1391–1398.
9. Hrycay EG, Bandiera SM: **Monooxygenase, peroxidase and peroxygenase properties and reaction mechanisms of cytochrome P450 enzymes.** In *Monooxygenase, Peroxidase and Peroxygenase Properties and Mechanisms of Cytochrome P450*. Edited by Hrycay EG, Bandiera SM.. Springer-Verlag Berlin; 2015: 60–80.
10. Ullrich R, Nüske J, Scheibner K, Spantzel J, Hofrichter M: **Novel haloperoxidase from the agaric basidiomycete Agrocybe aegerita oxidizes aryl alcohols and aldehydes.** *Appl Environ Microbiol* 2004, **70**:4575–4581.
11. Piontek K, Strittmatter E, Ullrich R, Gröbe G, Pecyna MJ, Kluge M, •• Scheibner K, Hofrichter M, Plattner DA: **Structural basis of substrate conversion in a new aromatic peroxygenase: P450 functionality with benefits.** *J Biol Chem* 2013, **288**:34767–34776.

This contribution reports the crystal structure of the peroxygenases from *Agrocybe aegerita*, which will put the basis for future protein engineering.

12. Kuhnel K, Blankenfeldt W, Terner J, Schlichting I: **Crystal structures of chloroperoxidase with its bound substrates and complexed with formate, acetate, and nitrate.** *J Biol Chem* 2006, **281**:23990–23998.
13. Gutierrez A, Babot ED, Ullrich R, Hofrichter M, Martinez AT, del Rio JC: **Regioselective oxygenation of fatty acids, fatty alcohols and other aliphatic compounds by a basidiomycete**

heme-thiolate peroxidase. *Arch Biochem Biophys* 2011, **514**:33-43.

14. Peter S, Kinne M, Wang XS, Ullrich R, Kayser G, Groves JT, Hofrichter M: **Selective hydroxylation of alkanes by an extracellular fungal peroxygenase.** *FEBS J* 2011, **278**:3667-3675.

15. Girhard M, Kunigk E, Tihovsky S, Shumyantseva VV, Urlacher VB: **Light-driven biocatalysis with cytochrome P450 peroxygenases.** *Biotechnol Appl Biochem* 2013, **60**:111-118.

16. Paul CE, Churakova E, Maurits E, Girhard M, Urlacher VB, Hollmann F: **In situ formation of H₂O₂ for P450 peroxygenases.** *Bioorg Med Chem* 2014, **22**:5692-5696.

17. Lucas F, Babot ED, Canellas M, del Rio JC, Kalum L, Ullrich R, Hofrichter M, Guallar V, Martinez AT, Gutierrez A: **Molecular determinants for selective C-25-hydroxylation of vitamins D-2 and D-3 by fungal peroxygenases.** *Catal Sci Technol* 2016, **6**:288-295.

This report nicely shows the differences in selectivity of two peroxygenases and proposes a structural explanation.

18. Peter S, Karich A, Ullrich R, Grobe G, Scheibner K, Hofrichter M: **Enzymatic one-pot conversion of cyclohexane into cyclohexanone: comparison of four fungal peroxygenases.** *J Mol Catal B Enzym* 2014, **103**:47-51.

19. Peter S, Kinne M, Ullrich R, Kayser G, Hofrichter M: **Epoxidation of linear, branched and cyclic alkenes catalyzed by unspecific peroxygenase.** *Enz Microb Technol* 2013, **52**:370-376.

20. Molina-Espeja P, Canellas M, Plou FJ, Hofrichter M, Lucas F, Guallar V, Alcalde M: **Synthesis of 1-naphthol by a natural peroxygenase engineered by directed evolution.** *ChemBioChem* 2016, **17**:341-349.

21. Kinne M, Poraj-Kobielska M, Ralph SA, Ullrich R, Hofrichter M, Hammel KE: **Oxidative cleavage of diverse ethers by an extracellular fungal peroxygenase.** *J Biol Chem* 2009, **284**:29343-29349.

22. Kluge M, Ullrich R, Scheibner K, Hofrichter M: **Stereoselective benzylic hydroxylation of alkylbenzenes and epoxidation of styrene derivatives catalyzed by the peroxygenase of *Agrocybe aegerita*.** *Green Chem* 2012, **14**:440-446.

23. Kinne M, Ullrich R, Hammel KE, Scheibner K, Hofrichter M: **Regioselective preparation of (R)-2-(4-hydroxyphenoxy)propionic acid with a fungal peroxygenase.** *Tetrahedron Lett* 2008, **49**:5950-5953.

24. Kinne M, Poraj-Kobielska M, Aranda E, Ullrich R, Hammel KE, Scheibner K, Hofrichter M: **Regioselective preparation of 5-hydroxypropranolol and 4'-hydroxydiclofenac with a fungal peroxygenase.** *Bioorg Med Chem Lett* 2009, **19**:3085-3087.

25. Kluge M, Ullrich R, Dolge C, Scheibner K, Hofrichter M: **Hydroxylation of naphthalene by aromatic peroxygenase from *Agrocybe aegerita* proceeds via oxygen transfer from H₂O₂ and intermediary epoxidation.** *Appl Microbiol Biotechnol* 2009, **81**:1071-1076.

26. Ullrich R, Hofrichter M: **Enzymatic hydroxylation of aromatic compounds.** *Cell Mol Life Sci* 2007, **64**:271-293.

27. Hollmann F, Arends IWCE: **Enzyme initiated radical polymerizations.** *Polymers* 2012, **4**:759-793.

28. Hu S, Hager LP: **Asymmetric epoxidation of functionalized cis-olefins catalyzed by chloroperoxidase.** *Tetrahedron Lett* 1999, **40**:1641-1644.

29. Aguilera S, Vazquez-Duhalt R, Tinoco R, Rivera M, Pecchi G, Alderete JB: **Stereoselective oxidation of R-(+)-limonene by chloroperoxidase from *Caldariomyces fumago*.** *Green Chem* 2008, **10**:647-653.

30. Rude MA, Baron TS, Brubaker S, Alibhai M, Del Cardayre SB, Schirmer A: **Terminal olefin (1-alkene) biosynthesis by a novel P450 fatty acid decarboxylase from *Jeotgalicoccus* species.** *Appl Environ Microbiol* 2011, **77**:1718-1727.

Transformation of carboxylic acids into terminal alkanes.

31. Dennig A, Kuhn M, Tassotti S, Thiessenhusen A, Gilch S, Bütler T, Haas T, Hall M, Faber K: **Oxidative decarboxylation of short-chain fatty acids to 1-alkenes.** *Angew Chem Int Ed* 2015, **54**:8819-8822.

32. Zachos I, Gassmeyer S, Bauer D, Sieber V, Hollmann F, Kourist R: **Photobiocatalytic decarboxylation for olefin synthesis.** *Chem Commun* 2015, **51**:1918-1921.

33. Wang J-B, Lonsdale R, Reetz MT: **Exploring substrate scope and stereoselectivity of P450 peroxygenase OleTJE in olefin-forming oxidative decarboxylation.** *Chem Commun* 2016.

34. Thiel D, Doknić D, Deska J: **Enzymatic aerobic ring rearrangement of optically active furylcarbinols.** *Nat Commun* 2014;5.

A very interesting extension of the reaction scope of peroxygenases.

35. Fernández-Fueyo E, Younes SHH, Rootselaar Sv, Aben RWM, Renirie R, Wever R, Holtmann D, Rutjes FPJT, Hollmann F: **A biocatalytic aza-Achmatowicz reaction.** *ACS Catal* 2016, **6**:5904-5907.

36. Getrey L, Krieg T, Hollmann F, Schrader J, Holtmann D: **Enzymatic halogenation of the phenolic monoterpenes thymol and carvacrol with chloroperoxidase.** *Green Chem* 2014, **16**:1104-1108.

37. Fernández-Fueyo E, van Wingerden M, Renirie R, Wever R, Ni Y, Holtmann D, Hollmann F: **Chemoenzymatic halogenation of phenols by using the haloperoxidase from *Curvularia inaequalis*.** *ChemCatChem* 2015, **7**:4035-4038.

This manuscript describes the application of an exceptionally (H₂O₂)-stable peroxygenase for chemical synthesis.

38. Carro J, Ferreira P, Rodriguez L, Prieto A, Serrano A, Balcells B, Arda A, Jimenez-Barbero J, Gutierrez A, Ullrich R et al.: **5-Hydroxymethylfurfural conversion by fungal aryl-alcohol oxidase and unspecific peroxygenase.** *FEBS J* 2015, **282**:3218-3229.

39. Wischand D, Hartung J: **Bromination of phenols in bromoperoxidase-catalyzed oxidations.** *Tetrahedron* 2012, **68**:9456-9463.

40. Wischand D, Radlow M, Hartung J: **Vanadate-dependent bromoperoxidases from *Ascophyllum nodosum* in the synthesis of brominated phenols and pyrroles.** *Dalton Trans* 2013, **42**:11926-11940.

41. Yaipakdee P, Robertson LW: **Enzymatic halogenation of flavanones and flavones.** *Phytochemistry* 2001, **57**:341-347.

42. Raugei S, Carloni P: **Structure and function of vanadium haloperoxidases.** *J Phys Chem B* 2006, **110**:3747-3758.

43. Kaup BA, Piantini U, Wust M, Schrader J: **Monoterpenes as novel substrates for oxidation and halo-hydroxylation with chloroperoxidase from *Caldariomyces fumago*.** *Appl Microbiol Biotechnol* 2007, **73**:1087-1096.

44. Roiban G-D, Reetz MT: **Expanding the toolbox of organic chemists: directed evolution of P450 monooxygenases as catalysts in regio- and stereoselective oxidative hydroxylation.** *Chem Commun* 2015, **51**:2208-2224.

45. Reetz MT: **Laboratory evolution of stereoselective enzymes: a prolific source of catalysts for asymmetric reactions.** *Angew Chem Int Ed* 2011, **50**:138-174.

46. Molina-Espeja P, Garcia-Ruiz E, Gonzalez-Perez D, Ullrich R, Alcalde M: **Directed evolution of unspecific peroxygenase from *Agrocybe aegerita*.** *Appl Environ Microbiol* 2014, **80**:3496-3507.

The first example of a directed evolution study of a peroxygenases.

47. Molina-Espeja P, Ma S, Mate DM, Ludwig R, Alcalde M: **Tandem yeast expression system for engineering and producing unspecific peroxygenase.** *Enz Microb Technol* 2015, **73**:74-29-33.

Improved fermentation yields of engineered peroxygenases.

48. Valderrama B, Ayala M, Vazquez-Duhalt R: **Suicide inactivation of peroxydases and the challenge of engineering more robust enzymes.** *Chem Biol* 2002, **9**:555-565.

49. Bankar SB, Bule MV, Singhal RS, Ananthanarayanan L: **Glucose oxidase — an overview.** *Biotechnol Adv* 2009, **27**:489-501.

50. Lutz S, Steckhan E, Liese A: **First asymmetric electroenzymatic oxidation catalyzed by a peroxidase.** *Electrochim Commun* 2004, **6**:583-587.

51. Krieg T, Hutmamn S, Mangold K-M, Schrader J, Holtmann D: **Gas diffusion electrode as novel reaction system for an electro-enzymatic process with chloroperoxidase.** *Green Chem* 2011, **13**:2686-2689.

52. Perez DI, Mifsud Grau M, Arends IWCE, Hollmann F: **Visible light-driven and chloroperoxidase-catalyzed oxygenation reactions.** *Chem Commun* 2009:6848-6850.

53. Churakova E, Kluge M, Ullrich R, Arends I, Hofrichter M, Hollmann F: **Specific photobiocatalytic oxyfunctionalization reactions.** *Angew Chem Int Ed* 2011, **50**:10716-10719.

54. Ni Y, Fernández-Fueyo E, Baraibar AG, Ullrich R, Hofrichter M, Yanase H, Alcalde M, van Berkel WJH, Hollmann F: **Peroxygenase-catalyzed oxyfunctionalization reactions promoted by the complete oxidation of methanol.** *Angew Chem Int Ed* 2016, **55**:798-801.
This contribution describes how methanol can be fully oxidized to CO₂ and provide a maximum of H₂O₂ equivalents to promote peroxygenase reactions.

55. Ni Y, Holtmann D, Hollmann F: **How green is biocatalysis? To calculate is to know.** *ChemCatChem* 2014, **6**:930-943.
Claiming environmental benignity is not enough. Estimating the environmental impact of a given reaction and comparing it to the state of the art is mandatory.

56. Tufvesson P, Lima-Ramos J, Nordblad M, Woodley JM: **Guidelines and cost analysis for catalyst production in biocatalytic processes.** *Org Proc Res Dev* 2010, **15**:266-274.
This article gives a realistic estimation about the real costs of an enzyme and thereby dispels common myths that enzymes are expensive.

57. Churakova E, Arends IWCE, Hollmann F: **Increasing the productivity of peroxidase-catalyzed oxyfunctionalization: a case study on the potential of two-liquid-phase systems.** *ChemCatChem* 2013, **5**:565-568.

58. Fernández-Fueyo E, Ni Y, Gomez Baraibar A, Alcalde M, van Langen LM, Hollmann F: **Towards preparative peroxygenase-catalyzed oxyfunctionalization reactions in organic media.** *J Mol Catal B Enzy* 2016.

59. Kille S, Zilly FE, Acevedo JP, Reetz MT: **Regio- and stereoselectivity of P450-catalysed hydroxylation of steroids controlled by laboratory evolution.** *Nat Chem* 2011, **3**:738-743.

60. Reetz MT: **Laboratory evolution of stereoselective enzymes as a means to expand the toolbox of organic chemists.** *Tetrahedron* 2012, **68**:7530-7548.

61. Wang JB, Reetz MT: **Biocatalysis — chiral cascades.** *Nat Chem* 2015, **7**:948-949.

62. Bornscheuer UT, Huisman GW, Kazlauskas RJ, Lutz S, Moore JC, Robins K: **Engineering the third wave of biocatalysis.** *Nature* 2012, **485**:185-194.