

Feedback between ice dynamics and bedrock deformation with 3D viscosity in Antarctica

van der Wal, W.; van Calcar, C.J.; de Boer, Bas; Blank, B.

DOI

[10.5194/egusphere-egu2020-19278](https://doi.org/10.5194/egusphere-egu2020-19278)

Publication date

2020

Document Version

Final published version

Citation (APA)

van der Wal, W., van Calcar, C. J., de Boer, B., & Blank, B. (2020). *Feedback between ice dynamics and bedrock deformation with 3D viscosity in Antarctica*. Abstract from EGU General Assembly 2020, Vienna, Austria. <https://doi.org/10.5194/egusphere-egu2020-19278>

Important note

To cite this publication, please use the final published version (if applicable). Please check the document version above.

Copyright

Other than for strictly personal use, it is not permitted to download, forward or distribute the text or part of it, without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license such as Creative Commons.

Takedown policy

Please contact us and provide details if you believe this document breaches copyrights. We will remove access to the work immediately and investigate your claim.



Feedback between ice dynamics and bedrock deformation with 3D viscosity in Antarctica

Wouter van der Wal^{1,2}, **Caroline van Calcar**², Bas de Boer³, and Bas Blank¹

¹Faculty of Aerospace Engineering, Delft University of Technology, Delft, Netherlands (w.vanderwal@tudelft.nl)

²Faculty of Civil Engineering and Geosciences, Delft University of Technology, Delft, Netherlands

³Earth and Climate Cluster, Faculty of Science, Vrije Universiteit Amsterdam, Amsterdam, Netherlands

Over glacial-interglacial cycles, the evolution of an ice sheet is influenced by Glacial isostatic adjustment (GIA) via two negative feedback loops. Firstly, vertical bedrock deformation due to a changing ice load alters ice-sheet surface elevation. For example, an increasing ice load leads to a lower bedrock elevation that lowers ice-sheet surface elevation. This will increase surface melting of the ice sheet, following an increase of atmospheric temperature at lower elevations. Secondly, bedrock deformation will change the height of the grounding line of the ice sheet. For example, a lowering bedrock height following ice-sheet advance increases the melt due to ocean water that in turn leads to a retreat of the grounding line and a slow-down of ice-sheet advance.

GIA is mainly determined by the viscosity of the interior of the solid Earth which is radially and laterally varying. Underneath the Antarctic ice sheet, there are relatively low viscosities in West Antarctica and higher viscosities in East Antarctica, in turn affecting the response time of the above mentioned feedbacks. However, most ice-dynamical models do not consider the lateral variations of the viscosity in the GIA feedback loops when simulating the evolution of the Antarctic ice sheet. The method developed by Gomez et al. (2018) includes the feedback between GIA and ice-sheet evolution and alternates between simulations of the two models where each simulation covers the full time period. We presents a different method to couple ANICE, a 3-D ice-sheet model, to a 3-D GIA finite element model. In this method the model computations alternates between the ice-sheet and GIA model until convergence of the result occurs at each timestep. We simulate the evolution of the Antarctic ice sheet from 120 000 years ago to the present. The results of the coupled simulation will be discussed and compared to results of the uncoupled ice-sheet model (using an ELRA GIA model) and the method developed by Gomez et al. (2018).