

A Lagrangian study of globally emitted aviation NO_x and associated short-term O₃ radiative forcing effects

Maruhashi, J.; Grewe, V.; Frömming, Christine; Jöckel, Patrick; Dedoussi, I.C.

DOI

[10.5194/egusphere-egu22-6646](https://doi.org/10.5194/egusphere-egu22-6646)

Publication date

2022

Document Version

Final published version

Citation (APA)

Maruhashi, J., Grewe, V., Frömming, C., Jöckel, P., & Dedoussi, I. C. (2022). *A Lagrangian study of globally emitted aviation NO_x and associated short-term O₃ radiative forcing effects*. Abstract from EGU General Assembly 2022, Vienna, Austria. <https://doi.org/10.5194/egusphere-egu22-6646>

Important note

To cite this publication, please use the final published version (if applicable). Please check the document version above.

Copyright

Other than for strictly personal use, it is not permitted to download, forward or distribute the text or part of it, without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license such as Creative Commons.

Takedown policy

Please contact us and provide details if you believe this document breaches copyrights. We will remove access to the work immediately and investigate your claim.

EGU22-6646, updated on 11 May 2022

<https://doi.org/10.5194/egusphere-egu22-6646>

EGU General Assembly 2022

© Author(s) 2022. This work is distributed under the Creative Commons Attribution 4.0 License.



A Lagrangian study of globally emitted aviation NO_x and associated short-term O₃ radiative forcing effects

Jin Maruhashi¹, Volker Grewe^{1,2}, Christine Frömming², Patrick Jöckel², and Irene Dedoussi¹

¹Faculty of Aerospace Engineering, Section Aircraft Noise and Climate Effects, Delft University of Technology, Delft, the Netherlands

²Institut für Physik der Atmosphäre, Deutsches Zentrum für Luft- und Raumfahrt, Oberpfaffenhofen, Germany

The resilient growth of air travel demands a comprehensive understanding of the climate effects from aviation emissions. The current level of knowledge of the environmental repercussions of CO₂ emissions is considerably higher than that of non-CO₂ emissions, which includes nitrogen oxides (NO_x), sulfur oxides (SO_x), other aerosols like black carbon (BC), water vapor and contrails. Aircraft NO_x emissions not only possess a high degree of uncertainty because of the non-linearity of the NO_x – O₃ chemistry, but are also responsible for producing the second strongest net warming effect out of all non-CO₂ climate forcers from aviation, right after contrails [1]. This study employs global-scale simulations to characterize the transport patterns of nitrogen oxides and assess their climate effects across several regions (North America, South America, Africa, Eurasia and Australasia) from January to March and July to September in 2014. Radiative forcing effects from the short-term increase in O₃, which are triggered by NO_x emissions, are estimated. These emissions, which are introduced at a typical cruising altitude, are modelled as Lagrangian air parcels that are transported within the ECHAM5/MESy Atmospheric Chemistry (EMAC) model [2]. In order to summarize the dynamical and radiative forcing characteristics of more than 10,000 simulated trajectories, a clustering approach with an adapted distance metric is applied. The method itself is an unsupervised machine learning algorithm, called QuickBundles [3], that is most commonly used in the field of neuroscience. A strong seasonal dependence is found for the contribution of NO_x emissions to O₃. In terms of residence times, NO_x emitted in Northern regions resides mainly in the upper mid-latitudes while those initiated in the South remain mostly in the Tropics. Due to pronounced zonal jets, the location of emission does not necessarily correspond to the region that will be most affected, i.e., an emission starting in N. America in July will induce the greatest warming in Europe.

[1] Lee, D.S., Fahey, D.W., Skowron, A., Allen, M.R., Burkhardt, U., Chen, Q., Doherty, S.J., Freeman, S., Forster, P.M., Fuglestedt, J., Gettelman, A., De León, R.R., Lim, L.L., Lund, M.T., Millar, R.J., Owen, B., Penner, J.E., Pitari, G., Prather, M.J., Sausen, R., Wilcox, L.J.: The contribution of global aviation to anthropogenic climate forcing for 2000 to 2018, *Atmospheric Environment*, Volume 244, 2021, 117834, ISSN 1352-2310, <https://doi.org/10.1016/j.atmosenv.2020.117834>.

[2] Jöckel, P., Kerkweg, A., Pozzer, A., Sander, R., Tost, H., Riede, H., Baumgaertner, A., Gromov, S.,

Kern, B., Development cycle 2 of the Modular Earth Submodel System (MESSy2), *Geoscientific Model Development*, 3, 717-752, doi: 10.5194/gmd-3-717-2010, 2010.

[3] Garyfallidis, E., Brett, M., Correia, M. M., Williams, G. B., Nimmo-Smith, I. QuickBundles, a Method for Tractography Simplification. *Frontiers in neuroscience*, 6, 175. <https://doi.org/10.3389/fnins.2012.00175>, 2012.