

Towards a new surrogate model for predicting short-term NO_x-O₃ effects from aviation using Gaussian processes

Rao, P.V.; Dwight, R.P.; Singh, D.; Maruhashi, J.; Dedoussi, I.C.; Grewe, V.; Frömming, Christine

DOI

[10.5194/egusphere-egu23-4337](https://doi.org/10.5194/egusphere-egu23-4337)

Publication date

2023

Document Version

Final published version

Published in

EGU General Assembly 2023

Citation (APA)

Rao, P. V., Dwight, R. P., Singh, D., Maruhashi, J., Dedoussi, I. C., Grewe, V., & Frömming, C. (2023). Towards a new surrogate model for predicting short-term NO_x-O₃ effects from aviation using Gaussian processes. In *EGU General Assembly 2023* [EGU23-4337] <https://doi.org/10.5194/egusphere-egu23-4337>

Important note

To cite this publication, please use the final published version (if applicable). Please check the document version above.

Copyright

Other than for strictly personal use, it is not permitted to download, forward or distribute the text or part of it, without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license such as Creative Commons.

Takedown policy

Please contact us and provide details if you believe this document breaches copyrights. We will remove access to the work immediately and investigate your claim.

EGU23-4337, updated on 08 May 2023

<https://doi.org/10.5194/egusphere-egu23-4337>

EGU General Assembly 2023

© Author(s) 2023. This work is distributed under the Creative Commons Attribution 4.0 License.



Towards a new surrogate model for predicting short-term $\text{NO}_x\text{-O}_3$ effects from aviation using Gaussian processes

Pratik Rao¹, Richard Dwight², Deepali Singh², Jin Maruhashi¹, Irene Dedoussi¹, Volker Grewe^{1,3}, and Christine Frömming³

¹Delft University of Technology, Faculty of Aerospace Engineering, Section Aircraft Noise and Climate Effects, Delft, Netherlands (p.v.rao@tudelft.nl).

²Delft University of Technology, Faculty of Aerospace Engineering, Section Aerodynamics, Wind Energy, and Propulsion, Delft, Netherlands.

³German Aerospace Center (DLR), Institute of Atmospheric Physics, Oberpfaffenhofen, Germany.

While efforts have been made to curb CO_2 emissions from aviation, the more uncertain non- CO_2 effects that contribute about two-thirds to the warming in terms of radiative forcing (RF), still require attention. The most important non- CO_2 effects include persistent line-shaped contrails, contrail-induced cirrus clouds and nitrogen oxide (NO_x) emissions that alter the ozone (O_3) and methane (CH_4) concentrations, both of which are greenhouse gases, and the emission of water vapour (H_2O). The climate impact of these non- CO_2 effects depends on emission location and prevailing weather situation; thus, it can potentially be reduced by advantageous re-routing of flights using Climate Change Functions (CCFs), which are a measure for the climate effect of a locally confined aviation emission. CCFs are calculated using a modelling chain starting from the instantaneous RF (iRF) measured at the tropopause that results from aviation emissions. However, the iRF is a product of computationally intensive chemistry-climate model (EMAC) simulations and is currently restricted to a limited number of days and only to the North Atlantic Flight Corridor. This makes it impossible to run EMAC on an operational basis for global flight planning. A step in this direction lead to a surrogate model called algorithmic Climate Change Functions (aCCFs), derived by regressing CCFs (training data) against 2 or 3 local atmospheric variables at the time of emission (features) with simple regression techniques and are applicable only in parts of the Northern hemisphere. It was found that in the specific case of O_3 aCCFs, which provide a reasonable first estimate for the short-term impact of aviation NO_x on O_3 warming using temperature and geopotential as features, can be vastly improved [1]. There is aleatoric uncertainty in the full-order model (EMAC), stemming from unknown sources (missing features) and randomness in the known features, which can introduce heteroscedasticity in the data. Deterministic surrogates (e.g. aCCFs) only predict point estimates of the conditional average, thereby providing an incomplete picture of the stochastic response. Thus, the goal of this research is to build a new surrogate model for iRF, which is achieved by :

1. Expanding the geographical coverage of iRF (training data) by running EMAC simulations in more regions (North & South America, Eurasia, Africa and Australasia) at multiple cruise flight

altitudes,

2. Following an objective approach to selecting atmospheric variables (feature selection) and considering the importance of local as well as non-local effects,

3. Regressing the iRF against selected atmospheric variables using supervised machine learning techniques such as homoscedastic and heteroscedastic Gaussian process regression.

We present a new surrogate model that predicts iRF of aviation $\text{NO}_x\text{-O}_3$ effects on a regular basis with confidence levels, which not only improves our scientific understanding of $\text{NO}_x\text{-O}_3$ effects, but also increases the potential of global climate-optimised flight planning.

References

[1] Rao, P.; et al. Case Study for Testing the Validity of $\text{NO}_x\text{-Ozone}$ Algorithmic Climate Change Functions for Optimising Flight Trajectories. *Aerospace* **2022**, *9*, 231. <https://doi.org/10.3390/aerospace9050231>