

Breakthrough in the challenging P450-catalyzed chemoenzymatic synthesis of C14-functionalized steroids

Reetz, Manfred T.; Hollmann, Frank

DOI

[10.1007/s11426-023-1625-6](https://doi.org/10.1007/s11426-023-1625-6)

Publication date

2023

Document Version

Final published version

Published in

Science China Chemistry

Citation (APA)

Reetz, M. T., & Hollmann, F. (2023). Breakthrough in the challenging P450-catalyzed chemoenzymatic synthesis of C14-functionalized steroids. *Science China Chemistry*, 66(9), 2435-2436.
<https://doi.org/10.1007/s11426-023-1625-6>

Important note

To cite this publication, please use the final published version (if applicable).
Please check the document version above.

Copyright

Other than for strictly personal use, it is not permitted to download, forward or distribute the text or part of it, without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license such as Creative Commons.

Takedown policy

Please contact us and provide details if you believe this document breaches copyrights.
We will remove access to the work immediately and investigate your claim.

Green Open Access added to TU Delft Institutional Repository

'You share, we take care!' - Taverne project

<https://www.openaccess.nl/en/you-share-we-take-care>

Otherwise as indicated in the copyright section: the publisher is the copyright holder of this work and the author uses the Dutch legislation to make this work public.

Breakthrough in the challenging P450-catalyzed chemoenzymatic synthesis of C14-functionalized steroids

Manfred T. Reetz^{1,2*} & Frank Hollmann³

¹Max-Planck -Institut für Kohlenforschung, Kaiser-Wilhelm-Platz 1, 45480 Mülheim, Germany;

²Tianjin Institute of Industrial Biotechnology, Chinese Academy of Sciences, Tianjin 300308, China;

³Department of Biotechnology, Delft University, van der Maasweg 9, 2629 HZ Delft, Netherlands

Received April 20, 2023; accepted May 5, 2023; published online May 12, 2023

Citation: Reetz MT, Hollmann F. Breakthrough in the challenging P450-catalyzed chemoenzymatic synthesis of C14-functionalized steroids. *Sci China Chem*, 2023, 66, <https://doi.org/10.1007/s11426-023-1625-6>

Intensive past research has demonstrated that cytochrome P450 monooxygenases are involved in the biosynthesis of numerous natural products and in human metabolism of various pharmaceuticals [1]. This inspired synthetic organic and pharmaceutical chemists to test them as catalysts in the oxidative hydroxylation of steroids. Unfortunately, wildtype P450s generally proved to show low levels of regio- and diastereoselectivity as well as suboptimal activity. In order to solve these (and other) challenging problems associated with P450s [2], efficient methods in semi-rational directed evolution were developed [3]. Rather than utilizing random mutagenesis of the type epPCR, or DNA shuffling, such focused mutagenesis techniques as Combinatorial Active-site Saturation Test (CAST) and Iterative Saturation Mutagenesis (ISM) were conceived and implemented [4], and subsequently applied to P450-catalyzed oxidative hydroxylation of steroids. For example, planned C7- and C16-selectivities were achieved [5,6]. Today, improved versions

of CAST/ISM are considered to have rational rather than just semi-rational character [3].

Zhou, Qu and co-workers have taken a significant step forward in selective P450-catalyzed synthesis of C14-functionalized steroids recently [7]. This difficult problem was solved in two phases, the first involving a clever strategy based on discovering the proper P450 monooxygenase and performing mutagenesis using alanine scanning, which allowed for the identification of hot-spots in the enzyme. The second phase was designed to ensure regio- and stereoselectivity in favor of the steroidal position C14, made possible by applying CAST/ISM. A number of structurally different steroids reacted regio- and stereoselectively (Figure 1, top). A theoretical analysis revealed the origin of selectivity (Figure 1, bottom). In addition to oxidative hydroxylation, chemoenzymatic sequences led to variously functionalized steroids, which are also valuable for potential pharmaceutical applications [7].

*Corresponding author (email: reetz@mpi-muelheim.mpg.de)




Figure 1 P450-catalyzed synthesis of C14 α -OH steroids. Top: structurally different steroids and their respective product C14 α alcohols. Below: in silico model of CYP14A with substrate **2a**. (a) The binding cavity of CYP14A. Substrate **2a**, haem and the binding cavity are colored cyan, brown, and light blue, respectively. The four critical residues, as well as V213, are labelled by the corresponding amino acids. (b) Comparison of WT enzyme and I111L-V124W. The mutated residues and C14 of the substrate are colored pink. (c) Comparison of the WT enzyme and I111L-M115K. The mutated residues and position C14 of the substrate are colored brown and pink, respectively. Adapted with permission from Ref. [7], copyright by The Author(s) (2023).

Conflict of interest The authors declare no conflict of interest.

- Guengerich FP, Waterman MR, Egli M. *Trends Pharmacol Sci*, 2016, 37: 625–640
- Whitehouse CJC, Bell SG, Wong LL. *Chem Soc Rev*, 2012, 41: 1218–1260
- Qu G, Li A, Acevedo-Rocha CG, Sun Z, Reetz MT. *Angew Chem Int Ed*, 2020, 59: 13204–13231
- Reetz MT, Carballeira JD. *Nat Protoc*, 2007, 2: 891–903
- Li A, Acevedo-Rocha CG, D'Amore L, Chen J, Peng Y, Garcia-Borràs M, Gao C, Zhu J, Rickerby H, Osuna S, Zhou J, Reetz MT. *Angew Chem Int Ed*, 2020, 59: 12499–12505
- Acevedo-Rocha CG, Gamble CG, Lonsdale R, Li A, Nett N, Hoenenreich S, Lingnau JB, Wirtz C, Fares C, Hinrichs H, Deege A, Mulholland AJ, Nov Y, Leys D, McLean KJ, Munro AW, Reetz MT. *ACS Catal*, 2018, 8: 3395–3410
- Song F, Zheng M, Wang J, Liu H, Lin Z, Liu B, Deng Z, Cong H, Zhou Q, Qu X. *Nat Synth*, 2023, doi: 10.1038/s44160-023-00280-z