

Bayesian inference of subduction processes from observations before and after the 2011 Tohoku earthquake

Marsman, Celine; Vossepoel, Femke; D'Acquisto, Mario; van Dinther, Ylona; Van de Wiel, Lukas; Govers, Rob

Publication date

2024

Document Version

Final published version

Citation (APA)

Marsman, C., Vossepoel, F., D'Acquisto, M., van Dinther, Y., Van de Wiel, L., & Govers, R. (2024). *Bayesian inference of subduction processes from observations before and after the 2011 Tohoku earthquake*. 640-640. Abstract from 39th General Assembly of the European Seismological Commission (ESC2024), Corfu, Greece.

Important note


To cite this publication, please use the final published version (if applicable). Please check the document version above.

Copyright

Other than for strictly personal use, it is not permitted to download, forward or distribute the text or part of it, without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license such as Creative Commons.

Takedown policy

Please contact us and provide details if you believe this document breaches copyrights. We will remove access to the work immediately and investigate your claim.



ESC2024-S34-159

Bayesian inference of subduction processes from observations before and after the 2011 Tohoku earthquake

Celine Marsman¹, Femke Vossepoel², Mario D'Acquisto³, Ylona van Dinther¹, Lukas van de Wiel, Rob Govers^{1*}

¹*Utrecht University, Utrecht, The Netherlands*, ²*Delft University of Technology, Delft, The Netherlands*, ³*GFZ German Research Centre for Geosciences, Potsdam, Germany*

* r.govers@uu.nl

We seek to quantify bulk viscoelastic flow, afterslip, and locking, within a rheological framework that is consistent over the entire earthquake cycle. We address this using an ensemble smoother. We construct a 2D finite element seismic cycle model with a power-law rheology in the asthenosphere. A priori information, such as a realistic temperature field and a coseismic slip distribution, is integrated into the model. Model pre-stresses are initialized during repeated earthquake cycles wherein the accumulated slip deficit is released entirely. We tailor the last earthquake to match the observed co-seismic slip of the 2011 Tohoku earthquake. The heterogeneous rheology structure is derived from the temperature field and experimental flow laws. Additionally, we simulate afterslip using a thin viscoelastic shear zone. We focus on constraining power-law flow parameters for the asthenosphere and the shear zone.

We assimilate 3D GEONET GNSS displacement time series acquired before and after the 2011 Tohoku earthquake. Power-law viscosity parameters are successfully retrieved for all domains. The data require separate viscoelastic domains in the mantle wedge above and below ~50 km depth. The sub-slab asthenosphere has viscoelastic properties that are distinctly different from the mantle wedge. The trade-off between the power-law activation energy and water fugacity hinders their individual estimation. The wedge viscosity is $>10^{19}$ Pa·s during the interseismic phase. Postseismic afterslip and bulk viscoelastic relaxation can be individually resolved from the surface deformation data. Afterslip is substantial between 40-50 km depth and extends to 80 km depth. Bulk viscoelastic relaxation in the wedge concentrates above 150 km depth with viscosities $<10^{18}$ Pa·s. Landward motion of the near-trench region occurs during the early postseismic period without the need for a separate low-viscosity channel below the slab.

