

## Quantitative microbial risk assessment for an indoor swimming pool with chlorination compared to a UV-based treatment

Peters, Marjolein; Keuten, Maarten; de Kreuk, Merle; Vrouwenvelder, Hans; Rietveld, Luuk; Medema, Gertjan

### Publication date

2017

### Document Version

Accepted author manuscript

### Citation (APA)

Peters, M., Keuten, M., de Kreuk, M., Vrouwenvelder, H., Rietveld, L., & Medema, G. (2017). *Quantitative microbial risk assessment for an indoor swimming pool with chlorination compared to a UV-based treatment*. Abstract from 7th International Conference 2017 on Swimming Pool and Spa Waters, Kos Island, Greece.

### Important note

To cite this publication, please use the final published version (if applicable). Please check the document version above.

### Copyright

Other than for strictly personal use, it is not permitted to download, forward or distribute the text or part of it, without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license such as Creative Commons.

### Takedown policy

Please contact us and provide details if you believe this document breaches copyrights. We will remove access to the work immediately and investigate your claim.

## Quantitative microbial risk assessment for an indoor swimming pool with chlorination compared to a UV-based treatment

Marjolein Peters<sup>a</sup>, Maarten Keuten<sup>a,b</sup>, Merle de Kreuk<sup>a</sup>, Hans Vrouwenvelder<sup>c</sup>, Luuk Rietveld<sup>a</sup>, GertJan Medema<sup>a</sup>

<sup>a</sup> Delft University of Technology, Section Sanitary Engineering, Delft, The Netherlands

<sup>b</sup> Hellebrekers Technieken, Nunspeet, The Netherlands

<sup>c</sup> Delft University of Technology, Department of Biotechnology, Delft, The Netherlands

### Aims

Most swimming pools use residual disinfectants like chlorine for disinfection. The use of chlorine has several drawbacks: some waterborne-pathogens are chlorine resistant and disinfection by-products (DBPs) may be formed which are associated with various health risks. Therefore, an alternative treatment was developed which consists of biological sand filtration, ultra-filtration and UV-disinfection. The goal of this study was to compare the microbial risks for bathers in a UV-disinfected pool compared to a chlorinated pool with the use of a quantitative microbial risk assessment.

### Methods

In this microbial risk assessment, the microbial release was calculated from multiple factors such as pool content, number of simultaneous swimmers, duration and frequency of swimming, ingestion of pool water, the hygienic behaviour of swimmers and the actual release of microbial cells per swimmer. The concentration of faecal bacteria was calculated from shedding experiments and known pathogen concentrations in faeces. The Dutch illness probability (283/1000) was used to calculate the number of infected bathers, which were all assumed to shed  $10^8$  faecal pathogens per g faeces. The used reference pathogens were *Campylobacter jejuni*, *E. coli* O157:H7 and *Salmonella enterica*. The removal of pathogens by treatment with UV-disinfection was set to 5-log units, every 4 hours, while during chlorination these 5-log units are known to be achieved in 30 seconds. The dose-response relationship for *E. coli* and *S. enterica* was simulated with a beta-Poisson distribution and for *C. jejuni* a hypergeometric function was used. The yearly risk of infection was calculated separately for each of the bacterial pathogens and a normal range sensitivity analysis was done.

### Results

The average bacterial cell concentration during opening hours in a UV-based swimming pool were the highest for *C. jejuni* ( $3.1 \times 10^{-3}$  cells L<sup>-1</sup>) > *S. enterica* ( $9.5 \times 10^{-4}$  cells L<sup>-1</sup>) > *E. coli* ( $7.2 \times 10^{-4}$  cells L<sup>-1</sup>). These calculated pathogen concentrations were about 180 times higher than calculated pathogen concentrations in a chlorinated swimming pool in which the averaged concentration was  $4.0 \times 10^{-6}$  cells L<sup>-1</sup> for pathogenic *E. coli* cells. Based on the average pathogen concentration during opening hours, the yearly risk of infection was calculated to be  $9.8 \times 10^{-8}$  for the chlorinated swimming pool and  $1.8 \times 10^{-5}$  for the UV-based swimming pool treatment. The yearly risks of infection of a UV-based treated swimming pool were the highest for *C. jejuni* ( $1.7 \times 10^{-3}$ ) > *E. coli* ( $1.8 \times 10^{-5}$ ) > *S. enterica* ( $3.5 \times 10^{-7}$ ). The simulated yearly risk of infection was found to be most sensitive for the number of bathers releasing pathogens.

### Conclusions

The yearly risks of infection of *E. coli* and *S. enterica* in a UV-based treated swimming pool were lower than the drinking water guidelines ( $10^{-4}$ ), but for *C. jejuni* it was higher. For a more complete picture of the health risks, the effects of disinfection by-products should also be taken into account in future risk assessments, as is the effect of other pathogens like *Pseudomonas*, *Cryptosporidium* and *Giardia*. UV-based treatment might be a good alternative for chlorination for some specific types of swimming pools.