

DNA Sequence Is a Major Determinant of Tetrasome Dynamics

Ordu, Orkide; Lusser, Alexandra; Dekker, Nynke H.

DOI

[10.1016/j.bpj.2019.07.055](https://doi.org/10.1016/j.bpj.2019.07.055)

Publication date

2019

Document Version

Accepted author manuscript

Published in

Biophysical Journal

Citation (APA)

Ordu, O., Lusser, A., & Dekker, N. H. (2019). DNA Sequence Is a Major Determinant of Tetrasome Dynamics. *Biophysical Journal*, 117(11), 2217-2227. <https://doi.org/10.1016/j.bpj.2019.07.055>

Important note

To cite this publication, please use the final published version (if applicable).
Please check the document version above.

Copyright

Other than for strictly personal use, it is not permitted to download, forward or distribute the text or part of it, without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license such as Creative Commons.


Takedown policy

Please contact us and provide details if you believe this document breaches copyrights.
We will remove access to the work immediately and investigate your claim.

Journal Pre-proof

DNA sequence is a major determinant of tetrasome dynamics

Orkide Ordu, Alexandra Lusser, Nynke H. Dekker



PII: S0006-3495(19)30712-X

DOI: <https://doi.org/10.1016/j.bpj.2019.07.055>

Reference: BPJ 9822

To appear in: *Biophysical Journal*

Received Date: 6 May 2019

Accepted Date: 30 July 2019

Please cite this article as: Ordu O, Lusser A, Dekker NH, DNA sequence is a major determinant of tetrasome dynamics, *Biophysical Journal* (2019), doi: <https://doi.org/10.1016/j.bpj.2019.07.055>.

This is a PDF file of an article that has undergone enhancements after acceptance, such as the addition of a cover page and metadata, and formatting for readability, but it is not yet the definitive version of record. This version will undergo additional copyediting, typesetting and review before it is published in its final form, but we are providing this version to give early visibility of the article. Please note that, during the production process, errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

© 2019

DNA sequence is a major determinant of tetrasome dynamics

Orkide Ordu¹, Alexandra Lusser², and Nynke H. Dekker^{1,*}

¹Department of Bionanoscience, Kavli Institute of Nanoscience, Delft University of Technology, van der Maasweg 9, 2629 HZ Delft, The Netherlands

²Division of Molecular Biology, Biocenter, Medical University of Innsbruck, Innrain 80-82, 6020 Innsbruck, Austria

*To whom correspondence should be addressed. Tel: +31(0)152783219; Email: N.H.Dekker@tudelft.nl

ABSTRACT

Eukaryotic genomes are hierarchically organized into protein-DNA assemblies for compaction into the nucleus. Nucleosomes, with the (H3-H4)₂ tetrasome as a likely intermediate, are highly dynamic in nature by way of several different mechanisms. We have recently shown that tetrasomes spontaneously change the direction of their DNA wrapping between left- and right-handed conformations, which may prevent torque build-up in chromatin during active transcription or replication. DNA sequence has been shown to strongly affect nucleosome positioning throughout chromatin. It is not known, however, whether DNA sequence also impacts the dynamic properties of tetrasomes. To address this question, we examined tetrasomes assembled on a high-affinity DNA sequence using freely orbiting magnetic tweezers. In this context, we also studied the effects of mono- and divalent salts on the flipping dynamics. We found that neither DNA sequence nor altered buffer conditions affect overall tetrasome structure. In contrast, tetrasomes bound to high-affinity DNA sequences showed significantly altered flipping kinetics, predominantly via a reduction in the lifetime of the canonical state of left-handed wrapping. Increased mono- and divalent salt concentrations counteracted this behaviour. Thus, our study indicates that high-affinity DNA sequences impact not only the positioning of the nucleosome, but that they also endow the subnucleosomal tetrasome with enhanced conformational plasticity. This may provide a means to prevent histone loss upon exposure to torsional stress, thereby contributing to the integrity of chromatin at high-affinity sites.

STATEMENT OF SIGNIFICANCE

Canonical (H3-H4)₂ tetrasomes possess high conformational flexibility, as evidenced by their spontaneous flipping between states of left- and right-handed DNA wrapping. Here, we show that these conformational dynamics of tetrasomes cannot be described by a fixed set of rates over all conditions. Instead, an accurate description of their behavior must take into account details of their loading, in particular the underlying DNA sequence. *In vivo*, differences in tetrasome flexibility could be regulated by modifications of the histone core or the tetrasomal DNA, and as such constitute an intriguing, potentially adjustable mechanism for chromatin to accommodate the torsional stress generated by processes such as transcription and replication.

INTRODUCTION

The nucleosome is the basic complex of chromatin in eukaryotic cells (1-3). It comprises 147 base pairs (bp) of deoxyribonucleic acid (DNA) wrapped around a disk-shaped assembly of eight histone proteins in ~1.7 turns (4-6). The histone octamer consists of two copies of two types of heterodimers, one of which is formed by histones H3 and H4, and the other by histones H2A and H2B (7,8). During nucleosome formation, first the two H3-H4 dimers assemble onto the DNA to form a tetrasome, after which the binding of two H2A/H2B dimers completes the full nucleosome (9). The resulting compaction upon DNA wrapping limits the accessibility of the

genome for essential cellular processes, such as transcription, replication and repair. Hence, the positioning and stability of nucleosomes play a key role in gene regulation and cellular function.

Recent research has provided significant new insights into the structure and especially the dynamics of nucleosomes (10). For example, studies using single-molecule techniques have revealed the intrinsically dynamic nature of nucleosomes in the form of ‘breathing’, i.e. the transient un- and rewinding of the nucleosomal DNA ends (11-14). Nucleosomes were also found to ‘gap’ by transiently opening and closing the two turns of nucleosomal DNA along the direction of the superhelical axis (15). The structure, stability and dynamics of nucleosomes have furthermore been shown to be extensively regulated by post-translational modifications (16), chaperones (17), and ATP-dependent remodelers (18). Furthermore, nucleosome composition and dynamics can be altered by forces and torques generated by molecular motors that process the genome (19,20). Such external influences, as well as changes in the ambient conditions, can cause nucleosomes to reorganize into different (sub)structures (21). Specifically, tetrasomes, which wrap ~80 bp of DNA around the (H3-H4)₂ tetramer (**Figure 1A**), have been observed as stable intermediates in several studies (22-28). Similar to nucleosomes, tetrasomes predominantly adopt a left-handed DNA wrapping, but interestingly, right-handed DNA wrapping has also been identified (29-34). Recently, we showed that tetrasomes can spontaneously switch between these two conformations without unbinding from the DNA (35-37) (**Figure 2A**). This conformational flexibility could mitigate the build-up of torsional stress in chromatin during e.g. transcription or replication and thereby prevent the eviction of histones from the DNA.

DNA sequence has been suggested to play an important role in determining the structural and dynamic properties of nucleosomes (38). The formation of a nucleosome is significantly affected by the ability of the underlying DNA to bend sharply (39,40). This feature has not only been examined in numerous *in vitro* studies (41), but it has also been observed at a global level *in vivo* (42). For example, in yeast it was shown that ~50% of the *in vivo* nucleosome positions can be attributed to sequence-specific features. In particular, high-affinity nucleosome binding sequences were found to be enriched in intergenic regions where an integral chromatin structure is important, while low affinity sequences were most dominant in highly transcribed regions, such as rRNA and tRNA genes (42). High-affinity sequences were also detected in strongly regulated genes, and it was proposed that such sequences may favour rapid reassembly of nucleosomes promoting repression of these genes under unfavourable conditions (42).

While structure and dynamics of high-affinity nucleosomes have also been studied extensively by *in vitro* single-molecule assays (43-48), an interesting open question is whether association of histones with high-affinity DNA sequences affects the dynamics of subnucleosomal structures, such as (H3-H4)₂ tetrasomes, which can arise in the course of transcription or other torque-generating processes. In this work we have addressed this question by a single molecule approach. Using freely-orbiting magnetic tweezers (FOMT) (49), we systematically investigated the structure and handedness dynamics of single (H3.1-H4)₂ tetrasomes assembled onto the well-characterized high-affinity Widom 601 nucleosome binding sequence (50). In addition, we have examined the contribution of different buffers commonly employed in chromatin studies (51-53) to the dynamics of high-affinity tetrasomes. Our results revealed intriguing effects of DNA sequence on handedness flipping dynamics that can be counteracted by mono- and divalent salts. Thus, our findings provide conceptual evidence for a model in which chromatin regions containing high-affinity nucleosomes may be better protected than other regions from complete histone loss during torque-generating processes, such as transcription or replication, due to inherently increased handedness flipping dynamics of (H3-H4)₂ tetrasomes.

MATERIALS AND METHODS

Preparation of DNA molecules

Tetrasome assembly was performed on linear double-stranded DNA (dsDNA) fragments of 1.96 kilo-base pair (kbp) length containing a single 601-sequence of 147 bp at their center. The two ends of the DNA fragments

were ligated to digoxigenin-coated (Roche Diagnostics, Basel, Switzerland) or biotin-coated (Roche Diagnostics) double-stranded fragments (handle) of 643 bp length, respectively, for immobilization and tethering. The schematics, sequence, and preparation of the DNA molecules are described in **Supplementary Figure S1** and **Supplementary Table SI**. Further details on the handles have been described in Ref. (37).

Expression and purification of proteins

The recombinant canonical *Drosophila* histones H3.1-H4 and nucleosome assembly protein-1 (NAP1) chaperones were expressed and purified as described in Ref. (35) and Ref. (37). While *in vivo* NAP1 has been identified as a chaperone for histones H2A and H2B (54), *in vitro* it has been found to also interact with histones H3 and H4 and prevent histone aggregation (35-37,55-60).

Preparation of histones and tetrasome assembly

Individual tetrasomes were assembled and monitored in real time using magnetic tweezers (see below) in three buffers with varying compositions of core components employed in previous studies (35-37,51-53) (**Table I**). To allow for direct comparison to our previous study (35), the protein samples were prepared similarly, by incubating 51 nanomolar (nM) of an equimolar solution of H3.1-H4 histones – either without or with 192 nM NAP1 – on ice for 30 min. The incubation buffers were identical to the measurement buffers (**Table I**), except for buffer A. In that case, the incubation buffer contained 0.25% (w/v) Polyethylene Glycol (PEG) and 0.25% (w/v) Polyvinyl Alcohol (PVA), and 0.1% (w/v) Bovine Serum Albumin (BSA), as in our previous study (35). The incubated protein solutions were then diluted by at most 1:5000, and 100 μ l of the diluted solution was flushed into the flow cell to assemble a single tetrasome. About 60 s after assembly, 100-300 μ l of measurement buffer was flushed through the flow cell to remove free proteins and to prevent further assembly of tetrasomes.

The details of the sample and flow cell preparation for magnetic tweezers experiments have been described in Ref. (37). Where the concentrations and volumes of DNA and magnetic bead solutions used in this work differ, this is stated in **Supplementary Table SII**. Five (out of total $N_{exp,total}=20$) experiments were performed by resuming the measurement on a tetrasome that did not disassemble during a preceding experiment. Two such experiments involved an exchange between buffers B and C (**Table I**) with 300-500 μ l of the respective measurement buffer: In one of these two experiments, a measurement was resumed in buffer C on a tetrasome that had been assembled before in buffer B, and in the other the reverse procedure was employed.

Magnetic tweezers instrumentation

The assembly and dynamics of an individual tetrasome was monitored for up to 10 h by directly measuring the length and angular position, i.e. the linking number, of a single DNA molecule in each experiment using FOMT (49) (**Figure 1B**). The magnetic tweezers hardware is detailed in Ref. (37). All experiments were performed at an exerted force of 0.6-0.7 piconewton (pN) and at room temperature (22 °C).

Data analysis

The analysis of the acquired data was performed using custom-written scripts with built-in functions in Matlab (Mathworks, Natick, MA, USA). An improved version of the custom-written step-finder algorithm described in Ref. (61) was used to detect stepwise changes in the time traces of the DNA length and DNA linking number. Subsequently, these fits were analyzed to identify simultaneous steps in both DNA length and DNA linking number within a time window of 19.1 s (see below), resulting from the assembly or disassembly of a tetrasome (**Figure 1C**, **Supplementary Figure S4**). With this approach, 48% ($n=13$) of all such steps ($N=27$) were automatically identified, while the remaining 52% ($n=14$) were corrected manually to better match the data. The retrieved sizes of these simultaneous changes in DNA length ($N=27$) and DNA linking number ($N=27$) form the basic quantities that characterize tetrasomes structure (**Supplementary Tables SIII and SIV**). Step sizes were also compared to the mean spatial resolution, determined by averaging the standard deviations (STDs) of the time trace profiles in all experiments (**Figure 1D**, **Supplementary Figure S5**).

The dynamics of tetrasome structure in terms of handedness flipping (**Figure 2A**) was separately analyzed in the parts of the time traces with a stable baseline in DNA length and DNA linking number (**Figure 2B**, **Supplementary Figure S7**). The change in DNA linking number upon flipping was determined by fitting a double Gaussian function to the DNA linking number data ($N=23$; **Figure 2B**, **Supplementary Table SIX**, and **Supplementary Figure S7**) and calculating the difference between the mean values. The probability for a tetrasome to occupy one of the two states was calculated from the relative ratio of the corresponding peak areas of the two Gaussian fits (**Supplementary Table SXI**).

The handedness dynamics of single, flipping (H3.1-H4)₂ tetrasomes was further analyzed in terms of the times spent in the left- or right-handed state using another custom-written algorithm based on Ref. (62). A threshold zone was defined by 1 STDs from the mean values of the two Gaussian distributions about their average (**Figure 2B**, **Supplementary Figure S7**). The details of the dwell-time analysis have been described in Ref. (37). Here, the corresponding DNA linking number data of the time traces of sufficiently long duration (>1600 s) were smoothed by filtering over $N=330$ ($N=1910$) points, corresponding to a time average of $\tau_{short} = 3.3$ s ($\tau_{long} = 19.1$ s). These time scales were obtained from autocorrelation analysis of the bead's angular fluctuations, as described in Refs. (37,49) (**Supplementary Figure S2**). The mean dwell time in each state was then determined by combining all datasets obtained in identical buffer conditions and fitting these to an exponential function (**Table II**, **Figure 3A-C**, **Supplementary Table SXII**, and **Supplementary Figure S8A-C**). For direct comparison, we also performed dwell-time analysis with the same settings on the partial time traces ($N=6$) of one of our earlier experiments published in Ref. (35). In that experiment, we note that three tetrasomes were assembled, while mainly a single tetrasome was flipping between the two conformations. Since a second tetrasome only flipped with a relative occurrence of 2.8% and 0.7% in two of the partial time traces, all dwell times in the right-handed state were assigned to one flipping tetrasome, yielding mean dwell time of $\tau_{D,right} = 14$ (+/-1) s ($N=161$). Since tetrasomes flip independently of one another, the mean dwell time of a single tetrasome in the left-handed state was obtained by multiplying the mean dwell time from the exponential fit by three, yielding $\tau_{D,left} = 183$ (+15/-12) s ($N=159$).

Further details of the data analysis have been described in Ref. (37). The significance of the similarity between the structural quantities obtained in the different conditions was assessed by respective unpaired two-sample t-tests (**Supplementary Tables SV-SVIII** and **SX**). The results for the dwell times obtained in the different conditions were checked for similarity by respective Wilcoxon rank-sum tests (**Supplementary Tables SXIII-SXV**). Finally, we note that the times measured in this study are an upper boundary due to the finite bead response time. The errors reported for the mean values obtained in this study correspond to 1 STD from the underlying distributions, unless indicated otherwise. The errors of calculated quantities were computed by error propagation.

Construction of energy diagrams

To construct the schematic energy diagrams of the left- and right-handed states of tetrasomes (**Figure 3D**, **Supplementary Figure S8D**), we have computed the free energy difference between them and estimated the height of the barrier separating them. The free energy difference can be calculated from the ratio of the probabilities $p_{left} = p$ and $p_{right} = 1-p$ for a tetrasome to be in the left- and right-handed conformations, respectively, using $\Delta E_{peak\ areas} = -k_B T \ln((1-p)/p)$ (see **Data Analysis** above, and **Supplementary Table SXI**). It can alternatively be computed from the ratio of the dwell times according to $\Delta E_{dwell-times} = -k_B T \ln(\tau_{D,right}/\tau_{D,left})$. The obtained results are summarized in **Table II** and **Supplementary Table SXII**. While the results from both approaches agree within error, we consider the energy differences calculated from the dwell times as more reliable due to the larger sample size. To estimate the height of the energy barrier ΔG^* relative to the left- and right-handed states of tetrasomes, we compute $\Delta G^* = -k_B T \ln(k/k_0)$, where the transition rates $k_{l \rightarrow r}$ and $k_{r \rightarrow l}$ equal the inverse of the dwell times $\tau_{D,left}$ and $\tau_{D,right}$, respectively, and the transition rate between the two states at zero force k_0 is assumed to be symmetric and estimated at $\sim 10^7$ s⁻¹ (37).

RESULTS

DNA sequence and buffer conditions do not affect the structural properties of tetrasomes

To assess whether high-affinity DNA sequences impact the formation and structure of tetrasomes, we loaded (H3.1-H4)₂ tetramers onto individual DNA molecules containing a single centrally positioned high-affinity 601-sequence (DNA_{w/601}) known for its strong nucleosome-positioning capability (50) with the help of the histone chaperone NAP1. Using FOMT (49), tetrasome assembly was recorded in real time by tracking the length z and linking number Θ of the DNA. In this setting, a single immobilized DNA molecule is tethered to a magnetic bead that is subject to constant force applied by a cylindrical permanent magnet, allowing its free rotation in the (x,y)-plane (**Figure 1B**). This force was set to 0.6-0.7 pN for direct comparison to our previous study of (H3.1-H4)₂ tetrasomes assembled by histone chaperone NAP1 on DNA molecules of random sequence (DNA_{random}) (35).

As in our previous studies with DNA_{random} (35-37), flushing in pre-incubated histone/NAP1-complexes ($n_{exp,subset}=9$ out of total $N_{exp,total}=20$) induced a simultaneous step-like change in both the length z and linking number Θ of DNA_{w/601}, indicating the assembly of a single tetrasome (**Figure 1C**). The mean values of these changes (**Supplementary Table SIII**) are in excellent agreement with those previously obtained for NAP1-loaded (H3.1-H4)₂ tetrasomes on DNA_{random} (35). Also in agreement with previous observations (35-37) is the lack of observable interactions between NAP1 chaperones and DNA molecules in the absence of histones (**Supplementary Figure S3**). To probe the effects of ambient conditions on tetrasome structure and dynamics, we performed the experiments in three different buffers employed in prior studies (35-37,51-53) that mainly differed in the concentrations of mono- and divalent salts (**Table I**) and obtained highly similar results (**Supplementary Table SIII**). We also flushed in (H3.1-H4)₂ tetramers in the absence of NAP1 ($n_{exp,subset}=11$). Thereby, DNA_{w/601} also exhibited a simultaneous step-like change in both measured quantities (**Supplementary Figure S4**). The mean values of the changes in length z and linking number Θ (**Supplementary Table SIII**) correspond excellently to those we obtained for (H3.1-H4)₂ tetrasomes loaded by NAP1 onto DNA_{w/601}, or onto DNA_{random} (35). The similarity of these results, which were furthermore independent of buffer conditions, indicate the formation of proper complexes in all cases. This was unexpected, since we never observed tetrasome formation in the absence of NAP1 in previous studies using random DNA sequences (35-37). Thus, the specific features of the high-affinity 601-sequence are capable of inducing spontaneous chaperone-independent assembly of histone tetramers.

Under the highly diluted conditions required for the controlled assembly of single tetrasomes (**Materials and Methods**), we also observed disassembly of tetrasomes in 40% ($n_{exp,subset}=8$) of all experiments ($N_{exp,total}=20$) within 3364 ± 765 s (mean ± 1 standard error of the mean (SEM); **Supplementary Figure S6**). The absolute mean values of the disassembly-associated changes in DNA length z and linking number Θ (**Supplementary Table SIV**) are in excellent agreement with those obtained upon assembly of tetrasomes (**Supplementary Table SIII**). Because disassembly occurred for both NAP1-loaded and spontaneously loaded tetrasomes in all buffer conditions, we can exclude a destabilizing effect of NAP1 and/or specific buffer conditions.

Given the similarity of the absolute changes in DNA length z and linking number Θ in all experiments in different buffer conditions ($N_{exp,total}=20$) (as assessed by respective t-tests, **Supplementary Tables SV-SVIII**), all results were grouped together to yield improved statistics. The absolute changes in DNA length z and DNA linking number Θ upon assembly and disassembly of single tetrasomes ($N_{counts,total}=27$) yielded mean values of $\Delta z_{(dis)ass} = 23 \pm 5$ nm ($n_{counts,subset}=25$) and $\Delta \Theta_{(dis)ass} = 0.9 \pm 0.2$ turns ($n_{counts,subset}=26$), respectively (**Figure 1D**). The distributions of the changes upon assembly or disassembly alone are shown in **Supplementary Figure S5A,B**, respectively. These results are in excellent agreement with values reported in other studies involving tetrasomes (25,30-37). Overall, our findings indicate that the high-affinity 601-sequence fully supports the

assembly of (H3.1-H4)₂ tetramers into proper tetrasome complexes, irrespective of the presence or absence of NAP1, or of buffer conditions.

High-affinity tetrasomes exhibit spontaneous DNA handedness flipping

To investigate the time-dependent behavior of tetrasomes stably bound to DNA_{w/601}, the FOMT measurements were continued for several hours following assembly. Like stably bound tetrasomes on DNA_{random} (35-37), high-affinity tetrasomes exhibited spontaneous dynamics in the handedness of their DNA wrapping (**Figure 2A**) irrespective of the buffer conditions. While the length z of the DNA molecules remained constant, the linking number Θ spontaneously fluctuated (or flipped) between two distinct mean values (**Figure 2B**, **Supplementary Figure S7**). Based on the negative and positive signs of these values, we deduce that they correspond to conformations of left- and right-handed DNA wrapping, respectively. As observed with tetrasomes on DNA_{random} (35-37), tetrasomes on DNA_{w/601} most frequently adopted the left-handed conformation. The difference in mean DNA linking number between the two observed states quantifies the induced change in tetrasome structure. The values of this quantity obtained under different experimental conditions, including tetrasomes loaded in the absence of NAP1, are summarized in **Supplementary Table SIX**. Due to the similarity of their values (as assessed by respective t-tests, **Supplementary Table SX**), the results from all experiments were pooled, which yielded a mean change in DNA linking number Θ upon flipping of $\Delta\Theta_{\text{flipping}} = 1.6 \pm 0.2$ turns ($N_{\text{counts,total}}=23$) (**Figure 2C**). These values are in excellent agreement with previous studies (35-37). Overall, our findings indicate that DNA sequence, the presence or absence of NAP1 during assembly, or buffer conditions do not affect the structural rearrangement of tetrasomes associated with handedness flipping.

High-affinity DNA sequences increase the kinetics of tetrasome handedness flipping

We further examined the handedness dynamics of single tetrasomes in terms of the times spent in each of the two states of DNA wrapping (**Materials and Methods**). To specifically examine the role of DNA sequence, we focused these studies on NAP1-assembled tetrasomes, comparing the dynamics of NAP1-assembled (H3.1-H4)₂ tetramers on either DNA_{w/601} or DNA_{random}. The details of tetrasome handedness dynamics upon spontaneous, NAP1-independent assembly can be found in the **Supplementary Results and Discussion** and in **Supplementary Figure S8**. Our results indicate that the underlying DNA sequence significantly alters the kinetics of tetrasome handedness flipping (as assessed by respective Wilcoxon rank-sum tests, **Supplementary Tables SXIII-SXV**). This can be deduced from the observation that a tetrasome on DNA_{w/601} spent an average time of $\tau_{\text{left}} = 37$ s in the left-handed state (in standard buffer A, **Figure 3A**, left), which is 4.9 ± 0.5 fold shorter than a tetrasome assembled onto DNA_{random} in the same buffer ($\tau_{\text{left}} = 183$ s, **Table II**). The right-handed state was affected to a lesser extent, displaying a lifetime $\tau_{\text{right}} = 28$ s (in standard buffer A, **Figure 3A**, right), which is 2.0 ± 0.3 times longer than a tetrasome on DNA_{random} in the same buffer ($\tau_{\text{right}} = 14$ s, **Table II**). These results clearly show that tetrasomes on high-affinity DNA sequences have a substantially enhanced tendency to flip from the left- to the right-handed state compared to tetrasomes on random DNA sequences (**Supplementary Table SXIII**). Since the experimental conditions were otherwise identical, these effects must arise from the 601-sequence in DNA_{w/601}.

We can translate the information from dwell-time measurements into schematic energy diagrams depicting the left- and right-handed states of tetrasomes (**Materials and Methods**; **Table II** and **Figure 3D**). Notably, the free energy differences ΔE between the left- and right-handed conformations of tetrasomes are reduced ~ 9 -fold on DNA_{w/601} ($\Delta E = 0.3 \pm 0.1$ k_BT) relative to DNA_{random} ($\Delta E = 2.6 \pm 0.1$ k_BT), reflecting the shift of the equilibrium occupancy towards the right-handed conformation for NAP1-loaded tetrasomes on DNA_{w/601}. This results predominantly from the smaller estimated energy difference between the left-handed state and the transition state (at the top of the barrier) for tetrasomes loaded onto DNA_{w/601} ($\Delta G^*_{\text{left}} \sim 19.7$ k_BT) relative to DNA_{random} ($\Delta G^*_{\text{left}} \sim 21.3$ k_BT).

We next examined the flipping dynamics of tetrasomes on DNA_{w/601} in two different buffer conditions. In buffer B (containing a two-fold increased concentration of monovalent salt in the absence of crowding agents, **Table I**), tetrasomes dwelled in the left-handed state for $\tau_{\text{left}} = 62$ s (**Figure 3B**, left), 1.7 ± 0.1 times longer than in

standard buffer A ($\tau_{left} = 37$ s, **Table II; Figure 3A**, left). The average lifetimes in the right-handed conformation were affected only slightly by the change to buffer B (**Figure 3B**, right; **Table II, Supplementary Tables SXIV and SXV**). In buffer C, which contained additional magnesium chloride relative to buffer B (**Table I**), a tetrasome dwelled in the left-handed conformation for an even longer time of $\tau_{left} = 141$ s (**Figure 3C**, top), which is 2.3 ± 0.2 -fold longer than in buffer B ($\tau_{left} = 62$ s, **Table II; Figure 3B**, left). Again, the mean dwell time in the right-handed state showed essentially no change (**Figure 3C**, bottom; **Figure 3B**, right).

These findings can also be transferred into schematic energy diagrams (**Materials and Methods; Figure 3D**). Because the predominant effect of increased monovalent salt concentration in buffer B for tetrasomes on DNA_{w/601} is an increase in the lifetime of the left-handed conformation, the free energy difference between the two states increases to $\Delta E = 0.6 \pm 0.1$ k_BT (compared to $\Delta E = 0.3 \pm 0.1$ k_BT in buffer A; **Table II**). The estimated barrier height relative to the left-handed state is altered to $\Delta G^*_{left} \sim 20.2$ k_BT, while the barrier height relative to the right-handed state remains essentially unaffected. Similarly, because the higher concentration of divalent ions in buffer C caused a further extension of the lifetime of the left-handed conformation, the free energy difference between the two states correspondingly increases to $\Delta E = 1.5 \pm 0.1$ k_BT (**Table II**). Thus, the estimated barrier height relative to the left-handed state for tetrasomes on DNA_{w/601} is larger in buffer C ($\Delta G^*_{left} \sim 21.1$ k_BT) than in buffer B ($\Delta G^*_{left} \sim 20.2$ k_BT), while the barrier height relative to the right-handed state remains essentially unaffected. In summary, higher concentrations of monovalent salt in the absence of crowding agents as well as divalent salts attenuate the handedness flipping dynamics of tetrasomes assembled on the high-affinity Widom 601-sequence by increasing the dwell time in the left-handed state and, thus, reducing the tendency for tetrasomes to flip to the right-handed state.

DISCUSSION

A strong component determining the positioning and stability of nucleosomes throughout chromatin is the DNA sequence (63). For instance, nucleosome maps obtained from chromatin assembled *in vitro* on purified yeast genomic DNA showed remarkable similarity to nucleosome maps from native chromatin (64). The observation that specific periodic patterns of AA, TT, TA or GC dinucleotides favour DNA wrapping around the histone core led to the development of algorithms to predict nucleosome positions in different organisms (42,65,66). High-affinity nucleosome binding sequences have been found to be enriched at the transcriptional start site, in intragenic locations or over genes subject to strong regulation (42). In the present study, we have investigated to which extent such high-affinity sites affect the structure and dynamics of subnucleosomal (H3-H4)₂ tetrasomes. Our previous single-molecule experiments revealed that the left-handed wrapping of DNA around the histone octamer is predominantly static, while the absence of H2A-H2B dimers results in highly dynamic flipping of DNA handedness between the left- and a right-handed state (35-37). However, these studies were carried out on tetrasomes assembled onto random DNA sequences, and it remained unclear whether high-affinity sequences, exemplified by the well-characterized Widom 601-sequence (50), would have an impact on the dynamics of tetrasomes. Our assessment of the assembly and structural features of (H3-H4)₂ tetrasomes on DNA including the 601-sequence revealed that the key characteristics of assembled tetrasomes were very similar to those assembled onto random DNA (35-37). Interestingly, however, we observed that the 601-sequence also supported spontaneous, histone chaperone-independent formation of long-lived, stable tetrasomes. As this property strongly distinguishes DNA including the high-affinity 601-sequence from DNA of random sequence, this suggests that proper tetrasome assembly occurs at the 601-sequence under these conditions. Such spontaneous chaperone-independent *in situ* assembly of histone tetramers may facilitate sample preparation in future single-molecule experiments. In the following, we assume that a similar sequence-specificity holds for the chaperoned assembly of tetrasomes onto DNA including the 601-sequence.

In contrast to the overall structural similarity between tetrasomes assembled onto high-affinity versus random DNA sequences, our analyses revealed a striking difference with respect to handedness flipping dynamics. The presence of the 601-sequence considerably decreased the lifetime of the canonical left-handed state of high-

affinity tetrasomes, corresponding to a reduction in the free energy ΔE between the left- and right-handed conformations, which results in enhanced flipping frequency.

These results may at first sight seem surprising given the high propensity of the 601-sequence to support nucleosome/tetrasome formation relative to other DNA sequences (67), which presumably derives from a combination of enhanced binding to histones and increased bendability or twistability of the DNA (38,50). Indeed, our observation that histone tetramers assemble spontaneously onto DNA molecules with a 601-sequence, but not onto DNA molecules of random sequence, emphasizes the strong affinity of the sequence for histones. Yet what could be the reason for the increased flipping dynamics of high-affinity tetrasomes? A possible explanation of these findings is that the stronger binding of the histone tetramer to the DNA reduces the impact of thermal forces on the histone-DNA contacts, such as possible disruption (in analogy to nucleosome 'breathing' (11-14)), and instead transmits them to the H3-H3 interface. This could in turn enhance the rotation at the H3-H3 interface that is proposed to underlie tetrasome handedness flipping (30-37). In this model, high-affinity sequences might not only promote energetically favorable histone-DNA interactions, but may also endow tetrasomes with enhanced ability to accommodate torsional stress by flipping between left- and right-handed conformation. This feature may be particularly important during transcription, when tetrasomes can arise from nucleosomes by loss or removal of H2A/H2B dimers, and when torque can be generated by elongating RNA polymerase II (68-70). Although most transcription-linked torsional stress is removed by topoisomerases (71,72), high-affinity tetrasomes may provide extra protection against complete histone ejection at crucial locations in the genome by their inherently stronger ability to absorb this stress.

In the presence of higher concentrations of mono- and divalent salt and an absence of synthetic crowding agents, the left-handed states of NAP1-loaded tetrasomes on DNA molecules with a 601-sequence become longer lived. Hence, these conditions decrease the tendency of tetrasomes to adopt the right-handed conformation. We consider a direct influence of the synthetic crowding agents PEG and PVA on tetrasome dynamics to be unlikely, as studies show that the influence of PEG (73) or PVA (74) on the kinetics of biological systems is negligible at the low concentrations (below 1%) employed in our standard buffer. Thus, these changes must arise from the higher concentration of salts, which will reduce the strength of electrostatic interactions through screening. Even a two-fold increase in salt concentration as employed here can induce small but significant changes in the properties of DNA (e.g. elasticity (75-77), torsional stiffness (78), and supercoiling (79)) and proteins (80), and in the case of histones, reduce their affinity for DNA (81). In agreement with the model proposed above, such a generic reduction in affinity could account for the observed increase of the lifetime of the left-handed state for tetrasomes on DNA molecules with a 601-sequence, because it would shift the balance between thermally induced opening of DNA-histone contacts and dimer rotation at the H3-H3 interface towards the former. Upon further addition of divalent salt, which increases electrostatic screening to an even larger extent, the dwell time in the left-handed state of NAP1-loaded tetrasomes on DNA molecules with a 601-sequence becomes more pronounced, as one might expect by analogy with the above. These findings also illustrate that the interpretation of results obtained using nucleosome-positioning sequences must take into account the specific buffer conditions employed.

In summary, we provide conceptual evidence that DNA sequence impacts tetrasome handedness flipping dynamics, which may contribute to the mitigation of torsional stress generated by molecular motors without accompanying eviction of histones.

AUTHOR CONTRIBUTIONS

O.O., A.L., and N.H.D. designed the research; O.O. performed the research and analyzed the data; O.O., A.L., and N.H.D. discussed the results and wrote the paper.

ACKNOWLEDGMENTS

The authors thank Artur Kaczmarczyk for insightful discussions, the laboratory of John van Noort for providing the pGEM3Z-mono601 plasmid and several stock solutions, Theo van Laar for preparing the DNA constructs, and Jacob Kerssemakers for providing the improved step-finder algorithm. The authors also acknowledge financial support from the funding agencies mentioned below.

FUNDING

This work was funded from the European Research Council (ERC) with a Consolidator Grant DynGenome (No:312221) to N.H.D and by the Austrian Science Fund (FWF) P31377-B30 to A.L.

CONFLICT OF INTEREST

None declared.

SUPPLEMENTARY DATA

A Supplementary Information file accompanies this manuscript.

REFERENCES

- Kornberg, R.D. (1974) Chromatin structure: A repeating unit of histones and DNA. *Science*, **184**, 868-871.
- Olins, A.L. and Olins, D.E. (1974) Spheroid chromatin units (v bodies). *Science*, **183**, 330-332.
- Oudet, P., Gross-Bellard, M. and Chambon, P. (1975) Electron microscopic and biochemical evidence that chromatin structure is a repeating unit. *Cell*, **4**, 281-300.
- Richmond, T.J., Finch, J.T., Rushton, B., Rhodes, D. and Klug, A. (1984) Structure of the nucleosome core particle at 7 Å resolution. *Nature*, **311**, 532-537.
- Luger, K., Mäder, A.W., Richmond, R.K., Sargent, D.F. and Richmond, T.J. (1997) Crystal structure of the nucleosome core particle at 2.8 Å resolution. *Nature*, **389**, 251-260.
- Davey, C.A., Sargent, D.F., Luger, K., Maeder, A.W. and Richmond, T.J. (2002) Solvent mediated interactions in the structure of the nucleosome core particle at 1.9 Å resolution. *J. Mol. Biol.*, **319**, 1097-1113.
- Klug, A., Rhodes, D., Smith, J., Finch, J.T. and Thomas, J.O. (1980) A low resolution structure for the histone core of the nucleosome. *Nature*, **287**, 509-516.
- Arents, G., Burlingame, R.W., Wang, B.C., Love, W.E. and Moudrianakis, E.N. (1991) The nucleosomal core histone octamer at 3.1 Å resolution: Tripartite protein assembly and a left-handed superhelix. *Proc. Natl. Acad. Sci. U. S. A.*, **88**, 10148-10152.
- Polo, S.E. and Almouzni, G. (2006) Chromatin assembly: A basic recipe with various flavours. *Curr. Opin. Genet. Dev.*, **16**, 104-111.
- Ordu, O., Lusser, A. and Dekker, N.H. (2016) Recent insights from in vitro single-molecule studies into nucleosome structure and dynamics. *Biophys. Rev.*, **8**, 33-49.
- Li, G., Levitus, M., Bustamante, C. and Widom, J. (2005) Rapid spontaneous accessibility of nucleosomal DNA. *Nat. Struct. Mol. Biol.*, **12**, 46-53.
- Koopmans, W.J.A., Brehm, A., Logie, C., Schmidt, T. and van Noort, J. (2007) Single-pair FRET microscopy reveals mononucleosome dynamics. *J. Fluoresc.*, **17**, 785-795.
- Miyagi, A., Ando, T. and Lyubchenko, Y.L. (2011) Dynamics of nucleosomes assessed with time-lapse high-speed atomic force microscopy. *Biochemistry*, **50**, 7901-7908.
- Wei, S., Falk, S.J., Black, B.E. and Lee, T.H. (2015) A novel hybrid single molecule approach reveals spontaneous DNA motion in the nucleosome. *Nucleic Acids Res.*, **43**, e111.
- Ngo, T.T. and Ha, T. (2015) Nucleosomes undergo slow spontaneous gaping. *Nucleic Acids Res.*, **43**, 3964-3971.
- Bowman, G.D. and Poirier, M.G. (2015) Post-translational modifications of histones that influence nucleosome dynamics. *Chem. Rev.*, **115**, 2274-2295.
- Gurard-Levin, Z.A., Quivy, J.P. and Almouzni, G. (2014) Histone chaperones: Assisting histone traffic and nucleosome dynamics. *Annu. Rev. Biochem.*, **83**, 487-517.
- Clapier, C.R. and Cairns, B.R. (2009), *Annu. Rev. Biochem.* Annual Reviews, Palo Alto, Vol. 78, pp. 273-304.
- Lavelle, C., Praly, E., Bensimon, D., Le Cam, E. and Croquette, V. (2011) Nucleosome-remodelling machines and other molecular motors observed at the single-molecule level. *FEBS J.*, **278**, 3596-3607.
- Lavelle, C. (2014) Pack, unpack, bend, twist, pull, push: the physical side of gene expression. *Curr. Opin. Genet. Dev.*, **25**, 74-84.
- Andrews, A.J. and Luger, K. (2011) Nucleosome structure(s) and stability: Variations on a theme. *Annu. Rev. Biophys.*, **40**, 99-117.
- Sollner-Webb, B., Camerini-Otero, R.D. and Felsenfeld, G. (1976) Chromatin structure as probed by nucleases and proteases: Evidence for the central role of histones H3 and H4. *Cell*, **9**, 179-193.
- Eickbush, T.H. and Moudrianakis, E.N. (1978) Histone core complex - octamer assembled by two sets of protein-protein interactions. *Biochemistry*, **17**, 4955-4964.

24. Dong, F. and van Holde, K.E. (1991) Nucleosome positioning is determined by the (H3-H4)₂ tetramer. *Proc. Natl. Acad. Sci. U. S. A.*, **88**, 10596-10600.
25. Brower-Toland, B.D., Smith, C.L., Yeh, R.C., Lis, J.T., Peterson, C.L. and Wang, M.D. (2002) Mechanical Disruption Of Individual Nucleosomes Reveals A Reversible Multistage Release Of DNA. *Proc. Natl. Acad. Sci. U. S. A.*, **99**, 1960-1965.
26. Hall, M.A., Shundrovsky, A., Bai, L., Fulbright, R.M., Lis, J.T. and Wang, M.D. (2009) High-resolution dynamic mapping of histone-DNA interactions in a nucleosome. *Nat. Struct. Mol. Biol.*, **16**, 124-129.
27. Böhm, V., Hieb, A.R., Andrews, A.J., Gansen, A., Rocker, A., Toth, K., Luger, K. and Langowski, J. (2011) Nucleosome accessibility governed by the dimer/tetramer interface. *Nucleic Acids Res.*, **39**, 3093-3102.
28. Sheinin, M.Y., Li, M., Soltani, M., Luger, K. and Wang, M.D. (2013) Torque modulates nucleosome stability and facilitates H2A/H2B dimer loss. *Nature Comm.*, **4**, 8.
29. Jackson, V. (1995) Preferential binding of histones H3 and H4 to highly positively coiled DNA. *Biochemistry*, **34**, 10607-10619.
30. Hamiche, A., Carot, V., Alilat, M., DeLucia, F., Odonohue, M.F., Revet, B. and Prunell, A. (1996) Interaction of the histone (H3-H4)₂ tetramer of the nucleosome with positively supercoiled DNA minicircles: Potential flipping of the protein from a left- to a right-handed superhelical form. *Proc. Natl. Acad. Sci. U. S. A.*, **93**, 7588-7593.
31. Hamiche, A. and Richard-Foy, H. (1998) The switch in the helical handedness of the histone (H3-H4)₂ tetramer within a nucleoprotein particle requires a reorientation of the H3-H3 interface. *J. Biol. Chem.*, **273**, 9261-9269.
32. Alilat, M., Sivolob, A., Revet, B. and Prunell, A. (1999) Nucleosome dynamics IV. Protein and DNA contributions in the chiral transition of the tetrasome, the histone (H3-H4)₂ tetramer-DNA particle. *J. Mol. Biol.*, **291**, 815-841.
33. Sivolob, A. and Prunell, A. (2000) Nucleosome Dynamics. V. Ethidium Bromide Versus Histone Tails In Modulating Ethidium Bromide-Driven Tetrasome Chiral Transition. A Fluorescence Study Of Tetrasomes On DNA Minicircles. *J. Mol. Biol.*, **295**, 41-53.
34. Sivolob, A., De Lucia, F., Alilat, M. and Prunell, A. (2000) Nucleosome Dynamics. VI. Histone Tail Regulation Of Tetrasome Chiral Transition. A Relaxation Study Of Tetrasomes On DNA Minicircles. *J. Mol. Biol.*, **295**, 55-69.
35. Vlijm, R., Lee, M., Lipfert, J., Lusser, A., Dekker, C. and Dekker, N.H. (2015) Nucleosome assembly dynamics involve spontaneous fluctuations in the handedness of tetrasomes. *Cell Rep.*, **10**, 216-225.
36. Vlijm, R., Lee, M., Ordu, O., Boltengagen, A., Lusser, A., Dekker, N.H. and Dekker, C. (2015) Comparing the assembly and handedness dynamics of (H3.3-H4)₂ tetrasomes to canonical tetrasomes. *PLoS One*, **10**, e0141267.
37. Ordu, O., Kremser, L., Lusser, A. and Dekker, N.H. (2018) Modification of the histone tetramer at the H3-H3 interface impacts tetrasome conformations and dynamics. *J. Chem. Phys.*, **148**, 123323.
38. Widom, J. (2001) Role of DNA sequence in nucleosome stability and dynamics. *Q. Rev. Biophys.*, **34**, 269-324.
39. Sekinger, E.A., Moqtaderi, Z. and Struhl, K. (2005) Intrinsic histone-DNA interactions and low nucleosome density are important for preferential accessibility of promoter regions in yeast. *Mol. Cell*, **18**, 735-748.
40. Anderson, J.D. and Widom, J. (2001) Poly(dA-dT) promoter elements increase the equilibrium accessibility of nucleosomal DNA target sites. *Mol. Cell Biol.*, **21**, 3830-3839.
41. Trifonov, E.N. and Nibhani, R. (2015) Review fifteen years of search for strong nucleosomes. *Biopolymers*, **103**, 432-437.
42. Segal, E., Fondufe-Mittendorf, Y., Chen, L., Thastrom, A., Field, Y., Moore, I.K., Wang, J.P. and Widom, J. (2006) A genomic code for nucleosome positioning. *Nature*, **442**, 772-778.
43. Blosser, T.R., Yang, J.G., Stone, M.D., Narlikar, G.J. and Zhuang, X. (2009) Dynamics of nucleosome remodelling by individual ACF complexes. *Nature*, **462**, 1022-1027.
44. Gansen, A., Valeri, A., Hauger, F., Felekyan, S., Kalinin, S., Toth, K., Langowski, J. and Seidel, C.A. (2009) Nucleosome disassembly intermediates characterized by single-molecule FRET. *Proc. Natl. Acad. Sci. U. S. A.*, **106**, 15308-15313.
45. Kruithof, M., Chien, F.T., Routh, A., Logie, C., Rhodes, D. and van Noort, J. (2009) Single-molecule force spectroscopy reveals a highly compliant helical folding for the 30-nm chromatin fiber. *Nat. Struct. Mol. Biol.*, **16**, 534-540.
46. Hodges, C., Bintu, L., Lubkowska, L., Kashlev, M. and Bustamante, C. (2009) Nucleosomal fluctuations govern the transcription dynamics of RNA polymerase II. *Science*, **325**, 626-628.
47. Bintu, L., Ishibashi, T., Dangkulwanich, M., Wu, Y.Y., Lubkowska, L., Kashlev, M. and Bustamante, C. (2012) Nucleosomal elements that control the topography of the barrier to transcription. *Cell*, **151**, 738-749.
48. Ngo, T.T., Zhang, Q., Zhou, R., Yodh, J.G. and Ha, T. (2015) Asymmetric unwrapping of nucleosomes under tension directed by DNA local flexibility. *Cell*, **160**, 1135-1144.
49. Lipfert, J., Wiggin, M., Kerssemakers, J.W.J., Pedaci, F. and Dekker, N.H. (2011) Freely orbiting magnetic tweezers to directly monitor changes in the twist of nucleic acids. *Nature Comm.*, **2**, 9.
50. Lowary, P.T. and Widom, J. (1998) New DNA sequence rules for high affinity binding to histone octamer and sequence-directed nucleosome positioning. *J. Mol. Biol.*, **276**, 19-42.
51. Meng, H., Andresen, K. and van Noort, J. (2015) Quantitative analysis of single-molecule force spectroscopy on folded chromatin fibers. *Nucleic Acids Res.*, **43**, 3578-3590.
52. Kaczmarczyk, A., Allahverdi, A., Brouwer, T.B., Nordenskiöld, L., Dekker, N.H. and van Noort, J. (2017) Single-molecule force spectroscopy on histone H4 tail cross-linked chromatin reveals fiber folding. *J. Biol. Chem.*, **292**, 17506-17513.
53. Kaczmarczyk, A., Brouwer, T.B., Pham, C., Dekker, N.H. and van Noort, J. (2018) In Lyubchenko, Y. L. (ed.), *Nanoscale Imaging: Methods and Protocols*. Springer New York, New York, NY, pp. 297-323.
54. Zlatanova, J., Seebart, C. and Tomschik, M. (2007) Nap1: taking a closer look at a juggler protein of extraordinary skills. *FASEB J.*, **21**, 1294-1310.
55. Vlijm, R., Smitshuijzen, J.S.J., Lusser, A. and Dekker, C. (2012) NAP1-assisted nucleosome assembly on DNA measured in real time by single-molecule magnetic tweezers. *PLoS One*, **7**, 11.
56. Fujii-Nakata, T., Ishimi, Y., Okuda, A. and Kikuchi, A. (1992) Functional analysis of nucleosome assembly protein, NAP-1. The negatively charged COOH-terminal region is not necessary for the intrinsic assembly activity. *J. Biol. Chem.*, **267**, 20980-20986.
57. Nakagawa, T., Bulger, M., Muramatsu, M. and Ito, T. (2001) Multistep Chromatin Assembly on Supercoiled Plasmid DNA by Nucleosome Assembly Protein-1 and ATP-utilizing Chromatin Assembly and Remodeling Factor. *J. Biol. Chem.*, **276**, 27384-27391.
58. Mazurkiewicz, J., Kepert, J.F. and Rippe, K. (2006) On the Mechanism of Nucleosome Assembly by Histone Chaperone NAP1. *J. Biol. Chem.*, **281**, 16462-16472.
59. Andrews, A.J., Chen, X., Zevin, A., Stargell, L.A. and Luger, K. (2010) The Histone Chaperone Nap1 Promotes Nucleosome Assembly by Eliminating Nonnucleosomal Histone DNA Interactions. *Mol. Cell*, **37**, 834-842.

60. McBryant, S.J., Park, Y.-J., Abernathy, S.M., Laybourn, P.J., Nyborg, J.K. and Luger, K. (2003) Preferential Binding of the Histone (H3-H4)₂ Tetramer by NAP1 Is Mediated by the Amino-terminal Histone Tails. *J. Biol. Chem.*, **278**, 44574-44583.
61. Kerssemakers, J.W.J., Munteanu, E.L., Laan, L., Noetzel, T.L., Janson, M.E. and Dogterom, M. (2006) Assembly dynamics of microtubules at molecular resolution. *Nature*, **442**, 709-712.
62. Lipfert, J., Skinner, G.M., Keegstra, J.M., Hensgens, T., Jager, T., Dulin, D., Köber, M., Yu, Z.B., Donkers, S.P., Chou, F.C. *et al.* (2014) Double-stranded RNA under force and torque: Similarities to and striking differences from double-stranded DNA. *Proc. Natl. Acad. Sci. U. S. A.*, **111**, 15408-15413.
63. Eslami-Mossallam, B., Schiessel, H. and van Noort, J. (2016) Nucleosome dynamics: Sequence matters. *Adv. Colloid Interface Sci.*, **232**, 101-113.
64. Kaplan, N., Moore, I.K., Fondufe-Mittendorf, Y., Gossett, A.J., Tillo, D., Field, Y., LeProust, E.M., Hughes, T.R., Lieb, J.D., Widom, J. *et al.* (2009) The DNA-encoded nucleosome organization of a eukaryotic genome. *Nature*, **458**, 362-366.
65. van der Heijden, T., van Vugt, J.J., Logie, C. and van Noort, J. (2012) Sequence-based prediction of single nucleosome positioning and genome-wide nucleosome occupancy. *Proc. Natl. Acad. Sci. U. S. A.*, **109**, E2514-2522.
66. Jiang, C. and Pugh, B.F. (2009) Nucleosome positioning and gene regulation: advances through genomics. *Nat. Rev. Genet.*, **10**, 161-172.
67. Thastrom, A., Lowary, P.T. and Widom, J. (2004) Measurement of histone-DNA interaction free energy in nucleosomes. *Methods*, **33**, 33-44.
68. Teves, S.S. and Henikoff, S. (2014) Transcription-generated torsional stress destabilizes nucleosomes. *Nat. Struct. Mol. Biol.*, **21**, 88-94.
69. Kouzine, F., Gupta, A., Baranello, L., Wojtowicz, D., Ben-Aissa, K., Liu, J., Przytycka, T.M. and Levens, D. (2013) Transcription-dependent dynamic supercoiling is a short-range genomic force. *Nat. Struct. Mol. Biol.*, **20**, 396-403.
70. Cole, H.A., Ocampo, J., Iben, J.R., Chereji, R.V. and Clark, D.J. (2014) Heavy transcription of yeast genes correlates with differential loss of histone H2B relative to H4 and queued RNA polymerases. *Nucleic Acids Res.*, **42**, 12512-12522.
71. Baranello, L., Wojtowicz, D., Cui, K., Devaiah, B.N., Chung, H.J., Chan-Salis, K.Y., Guha, R., Wilson, K., Zhang, X., Zhang, H. *et al.* (2016) RNA Polymerase II Regulates Topoisomerase 1 Activity to Favor Efficient Transcription. *Cell*, **165**, 357-371.
72. Levens, D., Baranello, L. and Kouzine, F. (2016) Controlling gene expression by DNA mechanics: emerging insights and challenges. *Biophys. Rev.*, **8**, 259-268.
73. Dupuis, N.F., Holmstrom, E.D. and Nesbitt, D.J. (2014) Molecular-crowding effects on single-molecule RNA folding/unfolding thermodynamics and kinetics. *Proc. Natl. Acad. Sci. U. S. A.*, **111**, 8464-8469.
74. Ariga, O., Sano, T. and Sano, Y. (1992) Effect of polyvinyl alcohol on kinetic and thermal properties of β -galactosidase. *J. Ferment. Bioeng.*, **74**, 120-122.
75. Baumann, C.G., Smith, S.B., Bloomfield, V.A. and Bustamante, C. (1997) Ionic effects on the elasticity of single DNA molecules. *Proceedings of the National Academy of Sciences*, **94**, 6185-6190.
76. Wenner, J.R., Williams, M.C., Rouzina, I. and Bloomfield, V.A. (2002) Salt Dependence of the Elasticity and Overstretching Transition of Single DNA Molecules. *Biophys. J.*, **82**, 3160-3169.
77. Manning, G.S. (2006) The Persistence Length of DNA Is Reached from the Persistence Length of Its Null Isomer through an Internal Electrostatic Stretching Force. *Biophys. J.*, **91**, 3607-3616.
78. Kriegel, F., Ermann, N., Forbes, R., Dulin, D., Dekker, N.H. and Lipfert, J. (2017) Probing the salt dependence of the torsional stiffness of DNA by multiplexed magnetic torque tweezers. *Nucleic Acids Res.*, **45**, 5920-5929.
79. Schlick, T., Li, B. and Olson, W.K. (1994) The influence of salt on the structure and energetics of supercoiled DNA. *Biophys. J.*, **67**, 2146-2166.
80. Damodaran, S. and Kinsella, J.E. (1981) The effects of neutral salts on the stability of macromolecules. A new approach using a protein-ligand binding system. *J. Biol. Chem.*, **256**, 3394-3398.
81. Rippe, K., Mazurkiewicz, J. and Kepper, N. (2008) In Dias, R. and Lindman, B. (eds.), *DNA Interactions with Polymers and Surfactants*. John Wiley & Sons, Inc.
82. Clapier, C.R., Chakravarthy, S., Petosa, C., Fernandez-Tornero, C., Luger, K. and Muller, C.W. (2008) Structure of the Drosophila nucleosome core particle highlights evolutionary constraints on the H2A-H2B histone dimer. *Proteins*, **71**, 1-7.

TABLES

Table I. Composition of the measurement buffers used in this work.

Measurement buffers	Buffer Components
A [Standard]	25 mM 4-(2-hydroxyethyl)-1-piperazineethane-sulfonic acid-KOH (HEPES-KOH (pH 7.5); Sigma-Aldrich, St. Louis, MO, United States), 0.1 mM ethylenediaminetetraacetic acid (EDTA; Sigma-Aldrich), 50 mM KCl (Merck, Darmstadt, Germany), 0.01% (w/v) BSA (Sigma-Aldrich), 0.025% (w/v) PEG (Sigma-Aldrich), 0.025% (w/v) PVA (Sigma-Aldrich)
B	10 mM HEPES (pH 7.5), 100 mM KCl, 0.2% (w/v) BSA, 0.1% (w/v) Tween-20 (Sigma-Aldrich), 10 mM NaN₃ (Merck)
C	10 mM HEPES (pH 7.5), 100 mM KCl, 0.2% (w/v) BSA, 0.1% (w/v) Tween-20, 10 mM NaN₃, 2 mM MgCl₂

Table II. Summary of quantified properties for NAP1-loaded tetrasomes. Specific buffers employed are indicated within brackets. The results for spontaneously loaded tetrasomes are provided in **Supplementary Table SXII**.

Quantity	(H3.1-H4) ₂ tetrasomes on DNA _{random}	(H3.1-H4) ₂ tetrasomes on DNA _{w/601}
$\Delta z_{(dis)ass}$ (nm)	24 (± 3) [*]	23 (± 5) ^x
$\Delta \theta_{(dis)ass}$ (turns)	0.73 (± 0.05) [*]	0.9 (± 0.2) ^x
$\Delta \theta_{flipping}$ (turns)	1.7 (± 0.1) [*]	1.6 (± 0.2) ^x
$\tau_{D,left}$ (s) with time average in filtering of 3.3 s [N=330]	183 (+15/-12) ⁺ [A]	37 (± 2) ^{**} [A] 62 (± 3) ^{**} [B] 141 (+11/-10) ^{**} [C]
$\tau_{D,left}$ (s) with time average in filtering of 19.1 s [N=1910]	417 (+60/-48) ⁺ [A]	67 (+5/-4) ^{**} [A] 121 (+8/-7) ^{**} [B] 231 (+24/-20) ^{**} [C]
$\tau_{D,right}$ (s) with time average in filtering of 3.3 s [N=330]	14 (± 1) [A]	28 (+2/-1) ^{**} [A] 33 (± 1) ^{**} [B] 30 (± 2) ^{**} [C]
$\tau_{D,right}$ (s) with time average in filtering of 19.1 s [N=1910]	34 (+5/-4) ^{**} [A]	48 (+4/-3) ^{**} [A] 64 (± 4) ^{**} [B] 44 (+5/-4) ^{**} [C]
$\tau_{D,left}/\tau_{D,right}$ with time average in filtering of 3.3 s [N=330]	13 (± 1) ⁺ [A]	1 (± 0) ⁺ [A] 2 (± 0) ⁺ [B] 4 (± 0) ⁺ [C]
$\tau_{D,left}/\tau_{D,right}$ with time average in filtering of 19.1 s [N=1910]	12 (± 3) ⁺ [A]	1 (± 0) ⁺ [A] 2 (± 0) ⁺ [B] 5 (± 1) ⁺ [C]
$\Delta E_{dwell-times}$ (k _B T)	2.6 (± 0.1) ⁺ [A]	0.3 (± 0.1) ⁺ [A] 0.6 (± 0.1) ⁺ [B] 1.5 (± 0.1) ⁺ [C]
$\Delta E_{peak-areas}$ (k _B T)	-	0.9 (± 0.7) ⁺ [A] 1.4 (± 0.8) ⁺ [B] 1.7 (\pm -----) [C]

*Values taken from Ref. (35)

^xValues obtained for spontaneously loaded tetrasomes on DNA_{w/601} are included for improved statistics

⁺Errors calculated by error propagation

^{**}Errors correspond to 68% confidence interval for estimated values from exponential fits


FIGURE CAPTIONS

Figure 1. Real-time assembly of single tetrasomes onto DNA with a 601-sequence. **A** Top view on about 80 bp of DNA (dark gray) bound to a tetrameric protein core consisting of the histones H3 (orange) and H4 (cyan). This image was created by modifying the structural data of the *Drosophila* nucleosome from the RCSB Protein Data Bank (PDB) with the PDB identification code 2PYO (82) using the PyMOL Molecular Graphics System, Version 1.8 Schrödinger, LLC. **B** Experimental assay based on FOMT (49). A single DNA construct (black) containing a 601-sequence (red) at its center ($\text{DNA}_{w/601}$) is attached to a coverslip (light blue) at one end, and tethered to a superparamagnetic bead (light brown) at the other end. Above the flow cell, a cylindrically shaped permanent magnet (dark blue/red), with its axis precisely aligned with the DNA tether, exerts a constant force (F) on the bead while allowing its free rotation in the (x,y)-plane (indicated by the black circular arrow). This enables the direct measurement of the DNA molecule's length z and linking number Θ , which upon the assembly of a tetrasome (orange) are changed by Δz and $\Delta\Theta$ (indicated by the red straight and circular arrows, respectively). Tetrasome assembly is induced by flushing in either $(\text{H3.1-H4})_2$ tetramers alone or pre-incubated histone/NAP1 complexes. Non-magnetic beads attached to the flow cell surface serve as reference for drift correction. This figure is adapted from Ref. (37) (<https://doi.org/10.1063/1.5009100>), with the permission of AIP Publishing. **C** Partial time traces of the length z (in nm, top panel) and the linking number Θ (in turns, bottom panel) of a $\text{DNA}_{w/601}$ molecule before and upon the assembly of a $(\text{H3.1-H4})_2$ tetrasome. The formation of a tetrasome simultaneously decreased both quantities in the form of a step identified using a step-finder algorithm (red lines) (**Materials and Methods**). About 60 s after assembly, free proteins were flushed out with measurement buffer (orange arrows) to prevent further histone binding. In this particular experiment, a tetrasome was assembled by flushing in NAP1/histone complexes (green arrows) in buffer A (**Table I**). A corresponding time trace of a $\text{DNA}_{w/601}$ molecule upon the assembly of a tetrasome from histone tetramers only is shown in **Supplementary Figure S4**. **D** Histogram of the changes in DNA length upon assembly and disassembly $\Delta z_{(dis)ass}$ (in nm, top panel) of single $(\text{H3.1-H4})_2$ tetrasomes ($N=27$) and in DNA linking number upon assembly and disassembly $\Delta\Theta_{(dis)ass}$ (in turns, bottom panel) of single $(\text{H3.1-H4})_2$ tetrasomes ($N=27$) in all buffers (**Table I**), shown together with the mean spatial resolution corresponding to the average 1 STD (16 nm and 0.5 turns; green lines) from all experiments. Data beyond the resolution limits (hatched area) was excluded from the determination of the mean change $\Delta z_{(dis)ass} = 23 \pm 5$ nm ($n=25$) and $\Delta\Theta_{(dis)ass} = 0.9 \pm 0.2$ turns ($n=26$).






Journal Pre-proof

Figure 2. The structural dynamics of single tetrasomes on DNA with a 601-sequence. **A** Representation of the two tetrasome conformations with the DNA wrapped in either a left-handed or a right-handed superhelix. Tetrasomes were observed to spontaneously flip between these two states (35-37). This figure is adapted from Ref. (37) (<https://doi.org/10.1063/1.5009100>), with the permission of AIP Publishing. **B** Partial time traces of the length z (in nm, top panel) and the linking number Θ (in turns, bottom panel) of a DNA molecule after the assembly of a (H3.1-H4)₂ tetrasome in buffer A (**Table I**) upon flushing in histone/NAP1 complexes. As indicated from the fit by the step-finder algorithm to the time trace (red line, left panel) and the fit of a mirrored gamma function (red line in histogram plot, right panel) to the skewed data, the DNA length remains constant. The DNA linking number spontaneously fluctuates, i.e. ‘flips’, between two states identified by fitting two Gaussian functions (white lines in histogram plot, right panel) underlying the full profile (red line in histogram plot, right panel). The two states correspond to a prevalent left-handed and a less adopted right-handed conformation of DNA wrapping, with the respective mean linking numbers $\Theta_{left} = -0.31 \pm 0.01$ turns and $\Theta_{right} = +1.38 \pm 0.06$ turns (dashed-dotted magenta lines, 95% confidence level for estimated values). Due to drift, the mean value for Θ_{left} obtained here is offset from the average change in DNA linking number upon tetrasome dis-/assembly (**Figure 1D**, bottom panel). The structural dynamics were quantified in terms of the dwell times in the two states based on a threshold zone (hatched orange area) that is bounded by 1 STD from each mean value (orange solid lines) about their average (solid magenta line) (**Materials and Methods**). A corresponding partial time trace of a DNA_{w/601} molecule upon the assembly of a tetrasome from histone tetramers only is shown in **Supplementary Figure S7**. **C** Histogram of the change in DNA linking number $\Delta\Theta_{flipping}$ (in turns) upon flipping of single (H3.1-H4)₂ tetrasomes in their handedness of DNA-wrapping in all buffers (**Table I**). The data yields a mean value of $\Delta\Theta_{flipping} = 1.6 \pm 0.2$ turns ($N=23$). The individual values are provided in **Supplementary Table SIX**.

Figure 3. Kinetics and energetics of single tetrasomes on DNA with a 601-sequence. **A** Histograms of the dwell times of a single NAP1-loaded (H3.1-H4)₂ tetrasome on a DNA_{w/601} molecule in the left- (left panel) and right-handed conformation (right panel) in buffer A (**Tables I and II**). Exponential fits (red lines) yielded a mean dwell time of $\tau_{D,left} = 37 \pm 2$ s ($N=371$) and $\tau_{D,right} = 28 \pm 2/-1$ s ($N=373$), respectively (68% confidence level for estimated values). **B** As in (A), but now in buffer B (**Tables I and II**). Exponential fits (red lines) yielded a mean dwell time of $\tau_{D,left} = 62 \pm 3$ s ($N=559$) and $\tau_{D,right} = 33 \pm 1$ s ($N=564$), respectively (68% confidence level for estimated values). **C** Histograms of the dwell times of a single NAP1-loaded (H3.1-H4)₂ tetrasome on a DNA_{w/601} molecule in the left-handed (top panel) and right-handed conformation (bottom panel) in buffer C (**Tables I and II**). Exponential fits (red lines) yielded a mean dwell time of $\tau_{D,left} = 141 \pm 11/-10$ s ($N=187$) and $\tau_{D,right} = 30 \pm 2$ s ($N=188$), respectively (68% confidence level for estimated values). All data in panels (A)-(C) were obtained by dwell-time analysis of the DNA linking number time traces filtered by averaging over 3.3 s ($N=330$) (**Materials and Methods**). **D** Schematic energy diagrams of single NAP1-loaded tetrasomes on DNA_{w/601} in all buffer conditions, based on the dwell-time values and the probabilities obtained from the linking number distributions (**Table II, Supplementary Tables SXI, SXIII-SXV**). The solid black lines illustrate the energy levels in buffer A (**Table I**) for a NAP1-loaded (H3.1-H4)₂ tetrasome on DNA_{random}, while the solid blue lines depict the energetics for a NAP1-loaded (H3.1-H4)₂ tetrasome on DNA_{w/601}. The green and orange dashed lines show the energy levels for a NAP1-loaded (H3.1-H4)₂ tetrasome on DNA_{w/601} in buffers B and C, respectively. The free energy differences (ΔE) between the left- and right-handed conformations of tetrasomes on DNA_{w/601}, with the respective energies E_{left} and E_{right} , are considerably decreased compared to tetrasomes loaded onto DNA_{random} (**Table II**). The heights of the energy barriers ΔG^*_{left} and ΔG^*_{right} are estimated from the rates $k_{l \rightarrow r}$ and $k_{r \rightarrow l}$, respectively. For tetrasomes on DNA_{w/601} relative to DNA_{random} in buffer A, ΔG^*_{left} decreases strongly, whereas ΔG^*_{right} is largely unchanged. In the presence of mono- and divalent salts in buffers B and C, tetrasomes on DNA_{w/601} become considerably longer-lived in the left-handed conformation (i.e. ΔG^*_{left} is higher) compared to tetrasomes on DNA_{w/601} in buffer A, while the right-handed state (hence ΔG^*_{right}) remains essentially unaltered.



A

Journal Pre-proof

B**C****D**

A

right-handed
DNA-wrapping

**B****C**