

Data-driven retrieval of primary plane-wave responses

Meles, Giovanni Angelo; Zhang, Lele; Thorbecke, Jan; Wapenaar, Kees; Slob, Evert

DOI

[10.5194/egusphere-egu2020-8606](https://doi.org/10.5194/egusphere-egu2020-8606)

Publication date

2020

Document Version

Final published version

Citation (APA)

Meles, G. A., Zhang, L., Thorbecke, J., Wapenaar, K., & Slob, E. (2020). *Data-driven retrieval of primary plane-wave responses*. Abstract from EGU General Assembly 2020, Vienna, Austria.
<https://doi.org/10.5194/egusphere-egu2020-8606>

Important note

To cite this publication, please use the final published version (if applicable).
Please check the document version above.

Copyright

Other than for strictly personal use, it is not permitted to download, forward or distribute the text or part of it, without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license such as Creative Commons.

Takedown policy

Please contact us and provide details if you believe this document breaches copyrights.
We will remove access to the work immediately and investigate your claim.



Data-driven synthesis of primary plane-wave responses

Giovanni Angelo Meles, Lele Zhang, Jan Thorbecke, Kees Wapenaar, and Evert Slob

Delft University of Technology, Faculty of Civil Engineering and Geosciences, The Hague, Netherlands (g.a.meles@tudelft.nl)

Seismic images provided by standard Reverse Time Migration are usually contaminated by artefacts associated with the migration of multiples.

Multiples can corrupt seismic images by producing both false negatives, i.e. by destructively interfering with primaries, and false positives, i.e. by focusing energy at unphysical interfaces. Free-surface multiples particularly affect seismic images resulting from marine data, while internal multiples strongly contaminate both land and marine data. Multiple prediction / primary synthesis methods are usually designed to operate on point source gathers, and can therefore be computationally demanding when large problems, involving hundreds of gathers, are considered.

In this contribution, a new scheme for fully data-driven retrieval of primary responses of plane-wave sources is presented. The proposed scheme, based on convolutions and cross-correlations of the reflection response with itself, extends a recently devised Marchenko point-sources primary retrieval method for to plane-wave source data. As a result, the presented algorithm allows fully data-driven synthesis of primary reflections associated with plane-wave source data. Once primary plane-wave responses are estimated, they are used for multiple-free imaging via standard reverse time migration. Numerical tests of increasing complexity demonstrate the potential of the proposed algorithm to produce multiple-free images only involving the migration of few datasets.

The plane-wave source primary synthesis algorithm discussed in this contribution could then be used as an initial and unexpensive processing step, potentially guiding more expensive target imaging techniques. Moreover, the method could be applied to large 3D problems for which standard methods are prohibitively expensive from a computational point of view.